

 $Head \ to \underline{www.savemyexams.com} \ for \ more \ awe some \ resources$

DP IB Maths: AI HL



4.10 Poisson Distribution

Contents

- * 4.10.1 Poisson Distribution
- * 4.10.2 Calculating Poisson Probabilities

4.10.1 Poisson Distribution

Your notes

Properties of Poisson Distribution

What is a Poisson distribution?

- A Poisson distribution is a discrete probability distribution
- A discrete random variable X follows a Poisson distribution if it counts the number of occurrences in a fixed time period given the following conditions:
 - Occurrences are independent
 - Occurrences occur at a uniform average rate for the time period (m)
- If X follows a Poisson distribution then it is denoted $X \sim \operatorname{Po}(m)$
 - m is the average rate of occurrences for the time period
- The formula for the probability of *r* occurrences is given by:


$$P(X=r) = \frac{e^{-m}m^r}{r!} \text{ for } r = 0,1,2,...$$

- e is Euler's constant 2.718...
- $r! = r \times (r-1) \times ... \times 2 \times 1$ and 0! = 1
- There is no upper bound for the number of occurrences
- You will be expected to use the distribution function on your GDC to calculate probabilities with the Poisson distribution

What are the important properties of a Poisson distribution?

- The **expected number (mean)** of occurrences is m
 - You are given this in the formula booklet
- The **variance** of the number of occurrences is *m*
 - You are given this in the formula booklet
 - Square root to get the standard deviation
- The **mean** and **variance** for a Poisson distribution are **equal**
- The distribution can be represented visually using a vertical line graph
 - The graphs have **tails to the right** for all values of *m*
 - As m gets larger the graph gets more symmetrical
- If $X \sim \operatorname{Po}(m)$ and $Y \sim \operatorname{Po}(\lambda)$ are independent then $X + Y \sim \operatorname{Po}(m + \lambda)$
 - This extends to *n* independent Poisson distributions $X_i \sim \text{Po}(m_i)$

$$X_1 + X_2 + \dots + X_n \sim Po(m_1 + m_2 + \dots + m_n)$$



2


0


4

6

8

10





Your notes



Head to www.savemyexams.com for more awesome resources

Modelling with Poisson Distribution

How do I set up a Poisson model?

- Identify what an occurrence is in the scenario
 - For example: a car passing a camera, a machine producing a faulty item
- Use proportion to find the mean number of occurrences for the given time period
 - For example: 10 cars in 5 minutes would be 120 cars in an hour if the Poisson model works for both time periods
- Make sure you **clearly state** what your **random variable** is
 - For example: let X be the number of cars passing a camera in 10 minutes

What can be modelled using a Poisson distribution?

- Anything that satisfies the two conditions
- For example, Let C be the number of calls that a helpline receives within a 15-minute period:

$C \sim Po(m)$

- An occurrence is the helpline receiving a call and can be considered independent
- The helpline receives calls at an average rate of m calls during a 15-minute period
- Sometimes a **measure of space** will be used instead of a time period
 - For example, the number of daisies that exist on a square metre of grass
- If the **mean** and **variance** of a discrete variable are **equal** then it might be possible to use a Poisson model

Examiner Tip

• An exam question might involve different types of distributions so make it clear which distribution is being used for each variable





Head to www.savemyexams.com for more awesome resources

Worked example

Jack uses Po(6.25) to model the number of emails he receives during his hour lunch break.

a) Write down two assumptions that Jack has made.

Jack has assumed that:

the emails that he receives are independent
he receives emails at a uniform average rate of
6.25 emails per hour during his lunch breaks

b) Calculate the standard deviation for the number of emails that Jack receives during his hour lunch breaks.

Formula	booklet	Poisson distribution $X \sim \text{Po}(m)$ Mean	E(X) = m
		Variance	Var(X) = m

$$\sigma^2 = 6.25$$
 $\sigma = \sqrt{6.25}$

Standard deviation = 2.5 emails



4.10.2 Calculating Poisson Probabilities

Your notes

Calculating Poisson Probabilities

Throughout this section we will use the random variable $X \sim \text{Po}(m)$. For a Poisson distribution X, the probability of X taking a non-integer or negative value is always zero. Therefore, any values mentioned in this section for X will be assumed to be non-negative integers. The value of M can be any real positive value.

How do I calculate P(X = x): the probability of a single value for a Poisson distribution?

- You should have a GDC that can calculate Poisson probabilities
- You want to use the "Poisson Probability Distribution" function
 - This is sometimes shortened to PPD, Poisson PD or Poisson Pdf
- You will need to enter:
 - The 'x' value the value of x for which you want to find P(X = x)
 - The 'λ' value the mean number of occurrences (m)
- Some calculators will give you the option of listing the probabilities for multiple values of x at once
- There is a formula that you can use but you are expected to be able to use the distribution function on your GDC

$$P(X=x) = \frac{e^{-m}m^x}{x!}$$

- where e is Euler's constant
- $X! = X \times (X 1) \times ... \times 2 \times 1 \text{ and } 0! = 1$

How do I calculate $P(a \le X \le b)$: the cumulative probabilities for a Poisson distribution?

- You should have a GDC that can calculate cumulative Poisson probabilities
 - Most calculators will find $P(a \le X \le b)$
 - Some calculators can only find $P(X \le b)$
 - The identities below will help in this case
- You should use the "Poisson Cumulative Distribution" function
 - This is sometimes shortened to PCD, Poisson CD or Poisson Cdf
- You will need to enter:
 - The lower value this is the value a
 - This can be zero in the case $P(X \le b)$
 - The upper value this is the **value b**
 - This can be a very large number (9999...) in the case $P(X \ge a)$
 - The ' λ ' value the mean number of occurrences (m)

How do I find probabilities if my GDC only calculates $P(X \le x)$?

• To calculate $P(X \le x)$ just enter x into the cumulative distribution function

- To calculate P(X < x) use:
 - $P(X < X) = P(X \le X 1)$ which works when X is a Poisson random variable
 - $P(X < 5) = P(X \le 4)$



- To calculate P(X > x) use:
 - $P(X > X) = 1 P(X \le X)$ which works for any random variable X
 - $P(X > 5) = 1 P(X \le 5)$
- To calculate P(X≥x) use:
 - $P(X \ge X) = 1 P(X \le X 1)$ which works when X is a Poisson random variable
 - $P(X \ge 5) = 1 P(X \le 4)$
- To calculate $P(a \le X \le b)$ use:
 - $P(a \le X \le b) = P(X \le b) P(X \le a 1)$ which works when X is a Poisson random variable
 - $P(5 \le X \le 9) = P(X \le 9) P(X \le 4)$

What if an inequality does not have the equals sign (strict inequality)?

■ For a Poisson distribution (as it is discrete) you could **rewrite all strict inequalities** (< and >) as **weak inequalities** (≤ and ≥) by using the identities for a Poisson distribution

$$P(X < X) = P(X \le X - 1) \text{ and } P(X > X) = P(X \ge X + 1)$$

- For example: $P(X < 5) = P(X \le 4)$ and $P(X > 5) = P(X \ge 6)$
- It helps to think about the range of integers you want
 - Identify the smallest and biggest integers in the range
- If your range has no minimum then use 0


•
$$P(X \le b) = P(0 \le X \le b)$$

■
$$P(a < X \le b) = P(a+1 \le X \le b)$$

- $P(5 < X \le 9) = P(6 \le X \le 9)$
- $P(a \le X < b) = P(a \le X \le b 1)$
 - $P(5 \le X < 9) = P(5 \le X \le 8)$
- $P(a < X < b) = P(a+1 \le X \le b-1)$
 - $P(5 < X < 9) = P(6 \le X \le 8)$

Worked example

The random variables $X \sim Po(6.25)$ and $Y \sim Po(4)$ are independent. Find:



$$m = 6.25$$
 (x) $x = 5$

$$P(X = 5) = 0.15341...$$

$$P(x=5) = 0.153 (3sf)$$

ii) $P(Y \le 5)$,

iii) P(X+Y>7).

Form the distribution
$$m = 6.25 + 4 = 10.25$$

$$P(X+Y>7) = 0.801 (3sf)$$

