

# DP IB Maths: AA HL



Your notes

## 3.8 Further Trigonometry

### Contents

- \* 3.8.1 Trigonometric Proof
- \* 3.8.2 Strategy for Trigonometric Equations



Your notes

## 3.8.1 Trigonometric Proof

### Trigonometric Proof

#### How do I prove new trigonometric identities?

- You can use trigonometric identities you already know to prove new identities
- Make sure you know how to find all of the trig identities in the formula booklet
  - The identity for tan, simple Pythagorean identity and the double angle identities for sin and cos are in the SL section
    - $\tan\theta = \frac{\sin\theta}{\cos\theta}$
    - $\cos^2\theta + \sin^2\theta = 1$
    - $\sin 2\theta = 2\sin\theta\cos\theta$
    - $\cos 2\theta = \cos^2\theta - \sin^2\theta = 2\cos^2\theta - 1 = 1 - 2\sin^2\theta$
- The reciprocal trigonometric identities for sec and cosec, further Pythagorean identities, compound angle identities and the double angle formula for tan
  - $\sec\theta = \frac{1}{\cos\theta}$
  - $\operatorname{cosec}\theta = \frac{1}{\sin\theta}$
  - $1 + \tan^2\theta = \sec^2\theta$
  - $1 + \cot^2\theta = \operatorname{cosec}^2\theta$
  - $\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$
  - $\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$
  - $\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$
  - $\tan 2\theta = \frac{2\tan\theta}{1 - \tan^2\theta}$
- The identity for cot is **not in the formula booklet**, you will need to remember it
  - $\cot\theta = \frac{1}{\tan\theta} = \frac{\cos\theta}{\sin\theta}$
- To prove an identity start on one side and proceed step by step until you get to the other side
  - It is more common to start on the left hand side but you can start a proof from either end
  - Occasionally it is easier to show that one side subtracted from the other is zero
  - You should not work on both sides simultaneously

#### What should I look out for when proving new trigonometric identities?

- Look for anything that could be a part of one of the above identities on either side
  - For example if you see  $\sin 2\theta$  you can replace it with  $2\sin\theta\cos\theta$
  - If you see  $2\sin\theta\cos\theta$  you can replace it with  $\sin 2\theta$
- Look for ways of reducing the number of different trigonometric functions there are within the identity
  - For example if the identity contains  $\tan\theta$ ,  $\cot\theta$  and  $\operatorname{cosec}\theta$  you could try
    - using the identities  $\tan\theta = 1/\cot\theta$  and  $1 + \cot^2\theta = \operatorname{cosec}^2\theta$  to write it all in terms of  $\cot\theta$
    - or rewriting it all in terms of  $\sin\theta$  and  $\cos\theta$  and simplifying
- Often you may need to trial a few different methods before finding the correct one
- Clever substitution into the **compound angle formulae** can be a useful tool for proving identities
  - For example rewriting  $\cos\frac{\theta}{2}$  as  $\cos\left(\theta - \frac{\theta}{2}\right)$  doesn't change the ratio but could make an identity easier to prove
- You will most likely need to be able to work with fractions and fractions-within-fractions
- Always keep an eye on the 'target' expression – this can help suggest what identities to use



Your notes

### Examiner Tip

- Don't forget that you can start a proof from either end – sometimes it might be easier to start from the left-hand side and sometimes it may be easier to start from the right-hand side
- Make sure you use the formula booklet as all of the relevant trigonometric identities are given to you
- Look out for special angles ( $0^\circ$ ,  $90^\circ$ , etc) as you may be able to quickly simplify or cancel parts of an expression (e.g.  $\cos 90^\circ = 0$ )



Your notes

### Worked example

Prove that  $8\cos^4\theta - 8\cos^2\theta + 1 = \cos 4\theta$ .

It is easiest to start on the right-hand side and apply the double angle formula for  $\cos 2\theta$ .

$$8\cos^4\theta - 8\cos^2\theta + 1 = \cos 4\theta$$

The form of the left-hand side suggests that the identity  $\cos 2A = 2\cos^2 A - 1$  would be more useful than the other options.

$$\begin{aligned}\cos 4\theta &= 2\cos^2 2\theta - 1 \\ &= 2(2\cos^2\theta - 1)^2 - 1 \\ &= 2(4\cos^4\theta - 4\cos^2\theta + 1) - 1 \\ &= 8\cos^4\theta - 8\cos^2\theta + 2 - 1\end{aligned}$$

$$\therefore \cos 4\theta = 8\cos^4\theta - 8\cos^2\theta + 1$$



Your notes

## 3.8.2 Strategy for Trigonometric Equations

### Strategy for Trigonometric Equations

#### How do I approach solving trig equations?


- You can solve trig equations in a variety of different ways
  - **Sketching a graph**
    - If you have your GDC it is always worth sketching the graph and using this to analyse its features
  - Using **trigonometric** identities, **Pythagorean** identities, the **compound** or **double angle** identities
    - Almost all of these are in the formula booklet, make sure you have it open at the right page
  - Using the **unit circle**
  - Factorising **quadratic** trig equations
    - Look out for quadratics such as  $5\tan^2x - 3\tan x - 4 = 0$
- The final rearranged equation you solve will involve **sin**, **cos** or **tan**
  - Don't try to solve an equation with **cosec**, **sec**, or **cot** directly

#### What should I look for when solving trig equations?

- Check the value of  $x$  or  $\theta$ 
  - If it is just  $x$  or  $\theta$  you can begin solving
  - If there are **different multiples** of  $x$  or  $\theta$  you will need to use the **double angle formulae** to get everything in terms of the same multiple of  $x$  or  $\theta$
  - If it is a **function** of  $x$  or  $\theta$ , e.g.  $2x - 15$ , you will need to **transform the range** first
    - You must remember to transform your solutions back again at the end
- Does it involve more than one trigonometric function?
  - If it does, try to **rearrange** everything to bring it to one side, you may need to **factorise**
  - If not, can you use an identity to reduce the number of different trigonometric functions?
    - You should be able to use identities to reduce everything to just one simple trig function (either  $\sin$ ,  $\cos$  or  $\tan$ )
- Is it **linear** or **quadratic**?
  - If it is linear you should be able to rearrange and solve it
  - If it is quadratic you may need to factorise first
    - You will most likely get two solutions, consider whether they both **exist**
    - Remember solutions to  $\sin x = k$  and  $\cos x = k$  only exist for  $-1 \leq k \leq 1$  whereas solutions to  $\tan x = k$  exist for all values of  $k$
- Are my solutions within the given range and do I need to find more solutions?
  - Be extra careful if your solutions are negative but the given range is positive only
  - Use a sketch of the graph or the unit circle to find the other solutions within the range
  - If you have a function of  $x$  or  $\theta$  make sure you are finding the solutions within the **transformed range**
    - Don't forget to transform the solutions back so that they are in the required range at the end




Your notes





Your notes







Your notes

ARE SOLUTIONS WITHIN GIVEN RANGE?  
DO I HAVE ALL SOLUTIONS?

YES

NO



CHECK WHETHER YOU HAVE FOUND ALL SOLUTIONS AND THAT THEY ARE WITHIN THE GIVEN RANGE  
(DON'T FORGET TO TRANSFORM YOUR SOLUTIONS BACK AT THE END IF THEY WERE FROM A TRANSFORMED RANGE)





Your notes

### Examiner Tip

- Try to use identities and formulas to reduce the equation into its simplest terms.
- Don't forget to check the function range and ensure you have included all possible solutions.
- If the question involves a function of  $x$  or  $\theta$  ensure you transform the range first (and ensure you transform your solutions back again at the end!).

### Worked example

Find the solutions of the equation  $(1 + \cot^2 2\theta)(5\cos^2 \theta - 1) = \cot^2 2\theta$  in the interval  $0 \leq \theta \leq 2\pi$ .

Move equivalent trig functions to the same sides:

$$\begin{aligned}
 5\cos^2 \theta - 1 &= \frac{\cot^2 2\theta}{1 + \cot^2 2\theta} && \text{divide both sides by } 1 + \cot^2 2\theta \\
 \cos 2\theta &= 2\cos^2 \theta - 1 && \\
 \therefore \cos^2 \theta &= \frac{1}{2} \cos 2\theta + \frac{1}{2} && \\
 5\left(\frac{1}{2} \cos 2\theta + \frac{1}{2}\right) - 1 &= \frac{\cot^2 2\theta}{\operatorname{cosec}^2 2\theta} && \\
 \frac{5}{2} \cos 2\theta + \frac{3}{2} &= \frac{\frac{\cos^2 2\theta}{\sin^2 2\theta}}{\frac{1}{\sin^2 2\theta}} && \begin{aligned} \cot \theta &= \frac{\cos \theta}{\sin \theta} \\ \operatorname{cosec} \theta &= \frac{1}{\sin \theta} \end{aligned} \\
 \frac{1}{2}(5\cos 2\theta + 3) &= \cos^2 2\theta && \text{Rearrange to form a} \\
 2\cos^2 2\theta - 5\cos 2\theta - 3 &= 0 && \text{quadratic in } \cos 2\theta \\
 (2\cos 2\theta + 1)(\cos 2\theta - 3) &= 0 && \\
 \cos 2\theta = -\frac{1}{2} \quad \text{or} \quad \cos 2\theta = 3 &&& \swarrow \text{no solutions}
 \end{aligned}$$

We are solving the equation for  $2\theta$  so we must transform the range first:  $0 \leq 2\theta \leq 4\pi$

$$2\theta = \cos^{-1}\left(-\frac{1}{2}\right) = \frac{2\pi}{3} \quad (\text{primary value})$$

$$\text{so } 2\theta = \frac{2\pi}{3}, \quad 2\pi - \frac{2\pi}{3} = \frac{4\pi}{3}, \quad \frac{2\pi}{3} + 2\pi = \frac{8\pi}{3}, \quad \frac{4\pi}{3} + 2\pi = \frac{10\pi}{3}$$

$$\theta = \frac{\pi}{3}, \frac{2\pi}{3}, \frac{4\pi}{3}, \frac{5\pi}{3}$$