

IB ⋅ SL ⋅ Biology

35 mins



Structured Questions

## **Cell Specialisation**

Stem Cells / Cell Specialisation

Total Marks	/35
Hard (2 questions)	/11
Medium (3 questions)	/17
Easy (3 questions)	

Scan here to return to the course

or visit savemyexams.com





## **Easy Questions**

1 (a)	Amoeba proteus is a unicellular organism that inhabits freshwater ponds and streams. At its largest size, Amoeba proteus can be visible to the naked eye.		
	State how the surface area to volume ratio changes as the <i>Amoeba proteus</i> grows.		
	(1 mark)		
(b)	Gaseous exchange allows the <i>Amoeba</i> to carry out aerobic respiration.		
	State which aspect of the surface area to volume ratio has the greater effect on the rate of gaseous exchange.		
	(1 mark)		
(c)	The contractile vacuole of the <i>Amoeba</i> helps to maintain its osmotic equilibrium by excreting excess water from the cell.		
	State which aspect of the surface area to volume ratio would depend on this organelle.		
	(1 mark)		
2	Glia and skeletal muscle cells are two examples of specialised cells that were once stem cells.		
	Define the term stem cell.		
	(2 marks)		



Embryonic stem cells are described as totipot	ent.
Define the term <i>totipotent</i> .	
	(2 marks



## **Medium Questions**

1 (a)	A patient with a spinal cord injury was treated using stem cell therapy using the following process:		
	<ol> <li>Stem cells were collected from the patient's own tissues</li> <li>Stem cells were cultured in laboratory for one month to increase their numbers</li> <li>Stem cells were injected into the injury site</li> <li>Stem cells were allowed to develop into nerve cells</li> </ol>		
	State <b>two</b> properties of stem cells that made this therapy possible.		
	(2 marks)		
(b)	Suggest why the stem cells used in part (a) were taken from the patient and not from a stem cell donor.		
	(2 marks)		

Caic	ulate the surface area to volume ratio of a cube where each edge measures 2 cm.
***************************************	(2 marks
	en first hatched, the young tadpoles of some frog species are less than 1.5 mm long have not yet developed gills.
	gest how these young tadpoles are able to get enough oxygen to their cells without eloped gills.
•••••	(2 marks
calcı	searcher calculated the surface area of a large number of frog eggs. She ulated the mean surface area to be 10.12 mm <sup>2</sup> . Frog eggs are generally spherical. calculated the surface area using the following equation: $4\pi r^2$ .
(i)	Rearrange this equation to find $r^2$ and use it to calculate the mean $r^2$ of these frog eggs.
(ii)	Calculate the mean diameter of these frog eggs. Give your answer to 3.s.f.
•••••	(3 marks)
Expl	ain why surface area to volume ratio is an important factor in limiting cell size.
	A re calco She (i)

(6 marks

## **Hard Questions**

1 (a)	shape is approximately cuboid.
	Calculate the surface area to volume ratio of this cell.
	(3 marks)
(b)	The palisade cell in part (a) has a large central vacuole. The vacuole has the same overall cuboid shape as the cell itself, but its outer surfaces are located 500 nm inside the plasma membrane of the cell.
	Calculate the surface area to cytoplasmic volume ratio.
	(3 marks)
(c)	Vacuoles have important roles in plant cell storage and support.
	Use your answers to parts (a) and (b) to suggest an additional role of vacuoles in plant cells.
	(2 marks)
2	Explain the importance of the surface area to volume ratio for a growing bacterial cell.

	(3 marks)