DP IB Chemistry: HL

7.1 Equilibrium

Contents

- ✤ 7.1.1 The State of Equilibrium
- ✤ 7.1.2 The Equilibrium Law
- * 7.1.3 Equilibrium Constant Relationships
- ★ 7.1.4 The Reaction Quotient
- ✤ 7.1.5 Le Chatelier's Principle
- ★ 7.1.6 Catalysts & Equilibrium





7.1.1 The State of Equilibrium

The State of Equilibrium

Reversible reaction

- Some reactions go to completion where the reactants are used up to form the products and the reaction stops when all of the reactants are used up
- In reversible reactions the products can react to reform the original reactants
- To show a reversible reaction, two half arrows are used: =






Dynamic equilibrium

- In a dynamic equilibrium the reactants and products are dynamic (they are constantly moving)
- In a dynamic equilibrium the **rate** of the **forward** reaction is the same as the rate of the **backward** reaction in a **closed system** and the **concentrations** of the **reactants** and **products** is **constant**
- There is no change in **macroscopic properties** such as colour and density as they depend on concentration




Page 2 of 20






The diagram shows a snapshot of a dynamic equilibrium in which molecules of hydrogen iodide are breaking down to hydrogen and iodine at the same rate as hydrogen and iodine molecules are reacting together to form hydrogen iodide



The diagram shows that the concentration of the reactants and products does not change anymore once equilibrium has been reached (equilibrium was approached using reactants)


Page 3 of 20

Your notes



The same equilibrium can be approached starting with the products

- A closed system is one in which none of the reactants or products escape from the reaction mixture
- In an **open system** some matter is lost to the surroundings
- When a reaction takes place entirely in solution, equilibrium can be reached in open flasks
- If the reaction involves gas, equilibrium can only be reached in a closed system



The diagram shows a closed system in which no carbon dioxide gas can escape and the calcium carbonate is in equilibrium with the calcium oxide and carbon dioxide

Page 4 of 20

Your notes

 $CaCO_3(s) \rightarrow CaO(s) + CO_2(g)$

The diagram shows an open system in which the calcium carbonate is continually decomposing as the carbon dioxide is lost causing the reaction to eventually go to completion

Examiner Tip

A common misconception is to think that the concentrations of the reactants and products is **equal**, however, they are **not** equal but **constant** (the concentrations are not changing). The dynamic equilibrium can be reached by starting either with the reactants or products. In both cases, the concentrations of the reactants and products remain constant once dynamic equilibrium has been reached.

7.1.2 The Equilibrium Law

The Equilibrium Constant

Equilibrium expression & constant

- The equilibrium expression is an expression that links the equilibrium constant, K_c, to the concentrations of reactants and products at equilibrium taking the stoichiometry of the equation into account
- So, for a given reaction:

 $aA + bB \Rightarrow cC + dD$

the K_c is expressed as follows:

$$K_{c} = \frac{[C]^{c}[D]^{d}}{[A]^{a}[B]^{b}}$$

 [A] AND [B] = EQUILIBRIUM REACTANT CONCENTRATIONS (mol dm⁻³)
[C] AND [D] = EQUILIBRIUM PRODUCT CONCENTRATIONS (mol dm⁻³)
a, b, c AND d = NUMBER OF MOLES OF REACTANTS AND PRODUCTS Copyright © Save My Exams. All Rights Reserved

Equilibrium expression linking the equilibrium concentration of reactants and products at equilibrium

- Solids are ignored in equilibrium expressions
- The K_c of a reaction is specific to a given equation

Worked example

Deduce the equilibrium constant expression for the following reactions

1. $Ag^+(aq) + Fe^{2+}(aq) = Ag(s) + Fe^{3+}(aq)$ 2. $N_2(g) + 3H_2(g) = 2NH_3(g)$ 3. $2SO_2(g) + O_2(g) = 2SO_3(g)$

Answer:



Page 6 of 20

Answer 1:

$$K_{c} = \frac{[Fe^{3+} (aq)]}{[Fe^{2+}(aq)] [Ag^{+}(aq)]}$$

[Ag(s)] is not included in the equilibrium expression as it is a solid

Answer 2:

$$K_{c} = \frac{[NH_{3}(g)]^{2}}{[N_{2}(g)][H_{2}(g)]^{3}}$$

Answer 3:

$$K_{c} = \frac{[SO_{3}(g)]^{2}}{[SO_{2}(g)]^{2} [O_{2}(g)]}$$

Your notes

Page 7 of 20

Fave MyExams Head to www.savemyexams.com for more awesome resources

The Size of Kc

• The size of K_c tells us how the equilibrium mixture is made up with respect to reactants and products



- If K_c > 1, the concentration of products is greater than the concentration of reactants and we say that the equilibrium lies to the right hand side
- If K_c < 1, then the concentration of reactants is greater than the concentration of products and we say that the equilibrium lies to the left hand side</p>
- K_c is a constant at a specified temperature
- Since temperature can affect the position of equilibrium it follows that K_c is dependent on temperature

Worked example

When the following reactions reach equilibrium, state whether the equilibrium mixture contains mostly reactants or products. Assume the value of K_c corresponds to the temperature of the reaction mixture

$$\begin{split} & 1. \ Ag^{+}(aq) + Fe^{2+}(aq) = Ag(s) + Fe^{3+}(aq) \quad K_{c} = 7.3 \times 10^{-26} \\ & 2. \ N_{2}(g) + 3H_{2}(g) = 2NH_{3}(g) \qquad K_{c} = 2.6 \times 10^{-18} \\ & 3. \ 2SO_{2}(g) + O_{2}(g) = 2SO_{3}(g) \qquad K_{c} = 5.0 \times 10^{13} \end{split}$$

Answer:

- 1 and 2: As K_c is very much smaller than 1 the denominator in the equilibrium expression must be much larger than the numerator so the concentration of the reactants is much larger than the concentration of products. The equilibrium mixture contains mostly reactants
- 3: As K_c is very much larger than 1 the numerator in the equilibrium expression must be much larger than the denominator so the concentration of the products is much larger than the concentration of reactants. The equilibrium mixture contains mostly products



7.1.3 Equilibrium Constant Relationships

Equilibrium Constant Relationships

- In the previous section we saw that the concentrations of the substances are raised to the **power of** the coefficients from the balanced equation
- This means the K_c expression is dependent on a specific equation
- For example, take the reaction between nitrogen and hydrogen to make ammonia

$$\frac{1}{2}N_{2(g)} + \frac{1}{2}H_{2(g)} \Rightarrow NH_{3(g)}$$

• The K_c expression for this reaction is:

$$K_{c} = \frac{[NH_{3}]}{[N_{2}]^{\frac{1}{2}}[H_{2}]^{\frac{3}{2}}}$$

If you double the stoichiometry the equation becomes

$$N_{2(q)} + 3H_{2(q)} \approx 2NH_{3(q)}$$

• The new **K**_c expression for this reaction is then:

NEW
$$K_c = \frac{[NH_3]^2}{[N_2][H_2]^3}$$

• What is the relationship between these two K_c values? You can probably see that when we double the coefficient the new K_c is the square of the original value:

NEW
$$K_c = \left(\text{ORIGINAL } K_c \right)^2$$

• If we reverse the equation:

$$2NH_{3(g)} = N_{2(g)} + 3H_{2(g)}$$

• K_c becomes the reciprocal of the original K_c value:

Page 9 of 20



F Save My Exams Head to <u>www.savemyexams.com</u> for more awesome resources

Your notes

REVERSE
$$K_c = \frac{[N_2][H_2]^3}{[NH_3]^2}$$

REVERSE $K_c = \frac{1}{ORIGINAL K_c}$

• Test your understanding in the following example:

Worked example

 K_c for 2NO₂(g) + F₂(g) = 2NO₂F(g) is 7.1 × 10³²What is K_c for the following reaction, at the same temperature?

 $NO_2F(g) \Rightarrow NO_2(g) + \frac{1}{2}F_2(g)$

A. 7.1 × 10³²

B.
$$\frac{1}{\sqrt{7.1 \times 10^{32}}}$$

C. $\frac{2}{7.1 \times 10^{32}}$
D. $\frac{1}{2 \times 7.1 \times 10^{32}}$

Answer:

The correct option is **B**.

 The original equation has been reversed and halved, so the K_c value must be square rooted and inverted to obtain the reciprocal

Examiner Tip

You must use square brackets in equilibrium constant expressions as they have a specific meaning, representing concentrations. In an exam answer you would lose the mark if you used round brackets

Page 10 of 20

7.1.4 The Reaction Quotient

The Reaction Quotient

- The **reaction quotient**, **Q**, is the ratio of products and reactants for a reaction that has **NOT** yet reached equilibrium
- The expression for **Q** is very similar to **K**_c:



- It is a useful concept because the size of Q can tell us how far a reaction is from equilibrium and in which direction the reaction proceeds
- For Example,
 - If Q = Kc then the reaction is at equilibrium, no net reaction occurs
 - If Q < Kc the reaction proceeds to the right in favour of the products
 - If **Q > Kc** the reaction **proceeds to the left** in favour of the reactants
- Using values of the concentrations of the substances present we can work out if a reaction is at equilibrium or not, as the following example shows:



Page 11 of 20

Worked example

The equilibrium constant for the following reaction:

 $COI_2(g) = CO(g) + I_2(g)$

is 5.1 x 10⁻² at 298 K

Deduce whether the following reaction mixture concentrations represent a reaction at equilibrium and for those not at equilibrium indicate the direction is proceeding:

Reaction mixture	[COl ₂ (g)]	[CO(g)]	[l ₂ (g)]
1	0.012	0.050	0.050
2	0.020	0.032	0.032
3	0.150	0.025	0.025
	Copyright © Save My Ex	ams. All Rights Reserved	

Answer:

The reaction quotient expression is

$$Q = \frac{[CO(g)][I_2(g)]}{[COI_2(g)]}$$

Reaction mixture 1:

$$Q = \frac{[0.050] [0.050]}{[0.012]} = 0.21$$

In this mixture $\mathbf{Q} >> \mathbf{K}_{\mathbf{c}}$, so Q has to decrease to reach $K_{\mathbf{c}}$. This means the reaction must be moving to the left, in order to reach equilibrium, so the reactants are favoured

Reaction mixture 2:

Page 12 of 20



$$Q = \frac{[0.032] [0.032]}{[0.020]} = 0.051$$

In this mixture, the value of **Q = Kc**, so the reaction is at equilibrium

Reaction mixture 3:

$$Q = \frac{[0.025] [0.025]}{[0.150]} = 0.0042$$

In this mixture $\mathbf{Q} < \mathbf{K}_{c}$, so Q has to increase to reach K_{c} . This means the reaction must be moving to the right, in order to reach equilibrium, so the products are favoured

Examiner Tip

The calculation of Q is not explicitly part of the SL course, just as calculating Kc values only comes in HL chemistry. However, a comparison of Q and Kc is relevant and the worked example is included only to illustrate how Q is determined from experimental data.



7.1.5 Le Chatelier's Principle

Le Chatelier's Principle

Position of the equilibrium

- The **position of the equilibrium** refers to the relative amounts of products and reactants in an equilibrium mixture.
- When the position of equilibrium shifts to the left, it means the concentration of reactants increases
- When the position of equilibrium shifts to the **right**, it means the concentration of **products** increases

Le Chatelier's principle

- Le Chatelier's principle says that if a change is made to a system at dynamic equilibrium, the position of the equilibrium moves to minimise this change
- The principle is used to predict changes to the position of equilibrium when there are changes in temperature, pressure or concentration

Effects of concentration

Effects of Concentration Table

CHANGE	HOW THE EQUILIBRIUM SHIFTS
INCREASE IN CONCENTRATION	EQUILIBRIUM SHIFTS TO THE RIGHT TO REDUCE THE EFFECT OF INCREASE IN THE CONCENTRATION OF A REACTANT
DECREASE IN CONCENTRATION	EQUILIBRIUM SHIFTS TO THE LEFT TO REDUCE THE EFFECT OF A DECREASE IN REACTANT (OR AN INCREASE IN THE CONCENTRATION OF PRODUCT)



Your notes



A. Using the reaction below:

 $CH_{3}COOH(I) + C_{2}H_{5}OH(I) \Rightarrow CH_{3}COOC_{2}H_{5}(I) + H_{2}O(I)$

Explain what happens to the position of equilibrium when:

1. More $CH_3COOC_2H_5(I)$ is added

- 2. Some $C_2H_5OH(I)$ is removed
- B. Use the reaction below:

 $Ce^{4+}(aq) + Fe^{2+}(aq) = Ce^{3+}(aq) + Fe^{3+}(aq)$

Explain what happens to the position of equilibrium when

3. Water is added to the equilibrium mixture

Answer:

Answer 1:

- The position of the equilibrium moves to the left and more ethanoic acid and ethanol are formed.
- The reaction moves in this direction to oppose the effect of added ethyl ethanoate, so the ethyl ethanoate decreases in concentration.

Answer 2:

- The position of the equilibrium moves to the left and more ethanoic acid and ethanol are formed.
- The reaction moves in this direction to oppose the removal of ethanol so more ethanol (and ethanoic acid) are formed from ethyl ethanoate and water.

Answer 3:

• There is no effect as the water dilutes all the ions equally so there is no change in the ratio of reactants to products.

Effects of pressure

• Changes in pressure only affect reactions where the reactants or products are gases

Effects of Pressure Table

CHANGE	HOW THE EQUILIBRIUM SHIFTS	
INCREASE IN PRESSURE	EQUILIBRIUM SHIFTS IN THE DIRECTION THAT PRODUCES THE SMALLER NUMBER OF MOLECULES OF GAS TO DECREASE THE PRESSURE AGAIN	
DECREASE IN PRESSURE	EQUILIBRIUM SHIFTS IN THE DIRECTION THAT PRODUCES THE LARGER NUMBER OF MOLECULES OF GAS TO INCREASE THE PRESSURE AGAIN	

Worked example

Predict the effect of increasing the pressure on the following reactions:

 $1. N_2O_4(g) \approx 2NO_2(g)$ 2. CaCO₃(s) \approx CaO(s) + CO₂(g)

Predict the effect of decreasing the pressure on the following reaction:

3. $2NO_2(g) = 2NO(g) + O_2(g)$

Answer:

Answer 1:

- The equilibrium shifts to the left as there are fewer gas molecules on the left.
- This causes a decrease in pressure.

Answer 2:

- The equilibrium shifts to the left as there are no gas molecules on the left but there is CO₂ on the right.
- This causes a decrease in pressure.

Answer 3:

- The equilibrium shifts to the right as there is a greater number of gas molecules on the right.
- This causes an increase in pressure.

Effects of temperature

Effects of Temperature Table

Page 16 of 20



Your notes

CHANGE	HOW THE EQUILIBRIUM SHIFTS
INCREASE IN TEMPERATURE	EQUILIBRIUM MOVES IN THE ENDOTHERMIC DIRECTION TO REVERSE THE CHANGE
DECREASE IN TEMPERATURE	EQUILIBRIUM MOVES IN THE EXOTHERMIC DIRECTION TO REVERSE THE CHANGE

Worked example

Using the reaction below:

 $H_2(g) + CO_2(g) = H_2O(g) + CO(g)$ $\Delta H = +41.2 \text{ kJ mol}^{-1}$

1. Predict the effect of increasing the temperature on this reaction

Using the reaction below:

 $Ag_2CO_3(s) \Rightarrow Ag_2O(s) + CO_2(g)$

2. Increasing the temperature increases the amount of $CO_2(g)$ at constant pressure. Is this reaction exothermic or endothermic?

Explain your answer

Answer:

Answer 1:

• The reaction will absorb the excess heat and since the forward reaction is endothermic, the equilibrium will shift to the right.

Answer 2:

• The reaction will absorb the excess heat and since this causes a shift of the equilibrium towards the right (as more CO₂(g) is formed) this means that the reaction is endothermic (because endothermic reactions favour the products).

Effects of catalysts

- A catalyst is a substance that increases the rate of a chemical reaction (they increase the rate of the **forward** and **reverse** reaction **equally**)
- Catalysts only cause a reaction to reach its equilibrium **faster**
- Catalysts therefore have **no effect** on the **position of the equilibrium** once this is reached

Page 17 of 20





You are not required to quote Le Chatelier's Principle in an exam, but you must know how to apply it to systems in equilibrium



7.1.6 Catalysts & Equilibrium

Catalysts & Equilibrium

Changes in concentration

- If all other conditions stay the same, the equilibrium constant K_c is **not affected** by any changes in concentration of the reactants or products
- For example, the decomposition of hydrogen iodide:

$$2HI \neq H_2 + I_2$$

The equilibrium expression is:

$$K_c = \frac{[H_2] [I_2]}{[HI]^2} = 6.25 \times 10^{-3}$$

Adding more HI makes the ratio of [products] to [reactants] smaller

To restore equilibrium, $[H_2]$ and $[I_2]$ increases and [HI] decreases

Equilibrium is restored when the ratio is 6.25×10^{-3} again

Changes in pressure

- A change in pressure only changes the position of the equilibrium (see Le Chatelier's principle)
- If all other conditions stay the same, the equilibrium constant K_c is **not affected** by any changes in pressure of the reactants and products

Changes in temperature

- Changes in temperature **change** the equilibrium constant K_c
- For an endothermic reaction such as:

$$2HI(g) = H_2(g) + I_2(g) \qquad K_c = \frac{[H_2][I_2]}{[HI]^2}$$

An increase in temperature:

[H₂] and [I₂] increases

[HI] decreases

Because [H₂] and [I₂] are **increasing** and [HI] is **decreasing**, the equilibrium constant K_c **increases**

For an exothermic reaction such as:

Page 19 of 20



SaveMyExams

$$2SO_2(g) + O_2(g) = 2SO_3(g)$$

$$K_c = \frac{[SO_3]^2}{[SO_2]^2[O_2]}$$



An increase in temperature:

 $[SO_3]\, \text{decreases}$

$[SO_2]$ and $[O_2]$ increases

Because [SO₃] decreases and [SO₂] and [O₂] increases the equilibrium constant K_c decreases

Presence of a catalyst

- If all other conditions stay the same, the equilibrium constant K_c is not affected by the presence of a catalyst
- A catalyst speeds up both the forward and reverse reactions at the same rate so the ratio of [products] to [reactants] remains unchanged

Worked example

An equilbrium is established in the following reaction:

AB(aq) + CD(aq) = AC(aq) + BD(aq) $\Delta H = +180 \text{ kJ mol}^{-1}$

Which factors would affect the value of in this equilibrium?

Answer:

- Only a change in temperature will affect the value of K_c and any other changes in conditions would result in the position of the equilibrium moving in such way to oppose this change.
- Adding a catalyst will increase the rate of reaction meaning the state of equilibrium will be reached faster but will have no effect on the position of the equilibrium and therefore K_c is unchanged.