

HLIB Physics



Gas Laws

Contents

- ***** Gas Pressure
- * Amount of Substance
- **★** Gas Laws
- ***** Ideal Gas Equation
- * Kinetic Theory of Gases
- * Derivation of the Kinetic Theory of Gases Equation
- * Average Kinetic Energy of a Molecule



Head to www.savemyexams.com for more awesome resources

Gas Pressure


Your notes

Calculating Gas Pressure

• Pressure is defined as


The force applied per unit area

- For example, when a drawing pin is pushed downwards:
 - It is pushed into the surface, rather than up towards the finger
 - This is because the sharp point is more **concentrated** (a small area) creating a **larger** pressure



When you push a drawing pin, it goes into the surface (rather than your finger)

- When an object is immersed in a liquid, the liquid will exert pressure, squeezing the object
 - The pressure exerted on objects in fluids creates **forces** against surfaces
 - These forces act at **90 degrees** (at right angles) to the surface






The pressure of a fluid on an object creates a force normal (at right angles) to the surface

• The equation for pressure is:

$$P = \frac{F}{A}$$

- Where:
 - P = pressure (Pa)
 - F = force(N)
 - A = cross-sectional area (m²)
- Pressure is measured in Pascals (Pa)
- This equation is only relevant when gas molecules exert a force **perpendicular** to the surface






Gas molecules bouncing off the walls of a container


- It is possible for someone to experience this force by closing their mouth and forcing air into their cheeks
- The strain on the cheeks is due to the force of the gas particles pushing at right angles to the cheeks
- This equation means:
 - If a force is spread over a large area it will result in a small pressure
 - If it is spread over a **small** area it will result in a **large** pressure











WEIGHT FROM HEELED SHOES IS SPREAD OVER A **SMALLER** AREA

THIS EXERTS A **HIGHER** PRESSURE ON THE GROUND

WEIGHT FROM FLAT SHOES IS SPREAD OVER A LARGER AREA

THIS EXERTS A **LOWER** PRESSURE ON THE GROUND


Copyright © Save My Exams. All Rights Reserved

High heels produce a higher pressure on the ground because of their smaller area, compared to flat shoes

Worked example

The diagram below shows the parts of the lifting machine used to move a platform of cross-sectional area $2.73 \times 10^{-2} \, \text{m}^2 \, \text{up}$ and down.





Copyright © Save My Exams. All Rights Reserved

The pump creates pressure in the liquid of 5.28×10^5 Pa to move the platform upwards.

Calculate the force that the liquid applies to the piston.

Answer:

Step 1: List the known quantities

- Cross-sectional area, $A = 2.73 \times 10^{-2} \,\mathrm{m}^2$
- Pressure, $P = 5.28 \times 10^5 \, \text{Pa}$

Step 2: Write down the relevant equation

$$p = \frac{F}{A}$$

Step 3: Rearrange for the force, F

$$F = p \times A$$



Step 4: Substitute the values into the equation

$$F = (5.28 \times 10^5) \times (2.73 \times 10^{-2}) = 14414.4$$

Step 5: Round to the appropriate number of significant figures and quote the correct unit

$$F = 14400 \text{ N} = 14.4 \text{ kN} (3 \text{ s.f})$$



Examiner Tip

Make sure A is always the **cross-sectional** area of the surface that the force is being applied upon, especially if there are multiple areas given in the question.

Amount of Substance

Your notes

Amount of Substance

- The **mole** is one of the seven SI base units
 - It is used to measure the **amount of substance**, not a mass
- One mole is defined as follows:

The amount of substance that contains as many elementary entities as the number of atoms in 12 g of carbon-12

- This amount of substance is exactly 6.02214076 x 10²³ elementary entities (i.e. particles, atoms, molecules)
 - At IB level, this number can be rounded to 6.02×10^{23}
- One mole of gas contains a number of particles (atoms or molecules) equal to the **Avogadro Constant**
- For example, 1 mole of sodium (Na) contains 6.02×10^{23} atoms of sodium
- The number of atoms can be determined if the number of moles is known by multiplying by N_A
 - For example: 2.0 mol of argon contains: $2.0 \times N_A = 2.0 \times 6.02 \times 10^{23} = 1.20 \times 10^{24}$ atoms
- The number of moles, n of a substance can be calculated using the equation

$$n = \frac{N}{N_A}$$

- Where:
 - N = number of particles (molecules or atoms, depending on the substance)
 - n = number of moles of gas (mol)
 - $N_A = 6.02 \times 10^{23} \,\text{mol}^{-1}$ (Avogadro constant)

Molar Mass

- One mole of any element is equal to the **relative atomic mass** of that element in grams
 - E.g. Helium has a relative atomic mass of 4 this means 1 mole of helium has a mass of 4 g
- If the substance is a compound, add up the relative atomic masses, for example, water (H₂O) is made up of
 - 2 hydrogen atoms (each with atomic mass of 1) and 1 oxygen atom (atomic mass of 16)
 - So, 1 mole of water would have a mass of $(2 \times 1) + 16 = 18$ g
- The **molar mass**, *m* of a substance is defined as the mass *m* of the substance divided by the amount (in moles) of that substance
- The molar mass is calculated as follows:

$$m_r = \frac{m}{n}$$

Where:

- $m_r = \text{molar mass in g mol}^{-1}$
- m = mass in grams (g)
- = n = number of moles (mol)



Worked example

120 moles of nitrogen gas are in a container.

Calculate the number of nitrogen gas molecules in the container.

Answer:

Step 1: List the known quantities

• Number of moles, n = 120

Step 2: Rearrange the number of moles equation for the number of molecules, N

$$N = nN_A$$

Step 3: Substitute in the values

$$N = (6.02 \times 10^{23}) \times 120 = 7.2 \times 10^{25}$$
 molecules

Worked example

A container is filled with 2.6×10^{20} molecules of argon gas which has a mass number of 40.

Calculate the total mass of argon in the container.

Answer:

Step 1: List the known quantities

- Number of argon gas molecules, $N = 2.6 \times 10^{20}$
- Molar mass of argon, $m_r = 40$ g mol⁻¹ (same as the atomic mass)

Step 2: Calculate the number of moles of argon in the container

$$n = \frac{N}{N_A} = \frac{2.6 \times 10^{20}}{6.02 \times 10^{23}} = 4.3 \times 10^{-4} \text{ moles}$$

Step 3: Calculate the total mass of the argon in the container

$$m = m_r \times n = 40 \times (4.3 \times 10^{-4}) = 0.0172 g$$





Examiner Tip

The value of the Avogadro constant is given on your data sheet, so you don't need to learn this value. However, it does help make calculations quicker if you do!

Take care with the units for molar mass, this is $\mathbf{g} \, \mathbf{mol}^{-1}$, so multiplying by the number of moles gives a mass in grams and **not** kilograms.



Gas Laws

Your notes

Gas Laws

• An ideal gas is one which obeys the relation:

- Where:
 - p = pressure of the gas (Pa)
 - $V = \text{volume of the gas } (m^3)$
 - T = thermodynamic temperature (K)
- Turning this into an equation gives:

$$\frac{PV}{T}$$
 = constant

• This is derived from the **empirical** gas laws for constant pressure, constant temperature and constant volume

Empirical Gas Laws

- The ideal gas laws are the experimental relationships between pressure P, volume V and the temperature T of an ideal gas
 - Boyle's Law (constant temperature)
 - Charles's Law (constant pressure)
 - Gay-Lussac's Law (constant volume)
- The mass and the number of molecules of the gas are assumed to be constant for all experiments

Boyle's Law

• If the temperature T of an ideal gas is constant, then **Boyle's Law** is given by:

$$P \propto \frac{1}{V}$$


- This means the pressure is **inversely proportional** to the volume of a gas
- The relationship between the pressure and volume for a fixed mass of gas at constant temperature can also be written as:

$$P_1 V_1 = P_2 V_2$$

- Where:
 - P₁ = initial pressure (Pa)
 - P_2 = final pressure (Pa)
 - V_1 = initial volume (m³)
 - V_2 = final volume (m³)

• The variation of volume and pressure at a constant temperature is shown below:






Copyright © Save My Exams, All Rights Reserved

Boyle's Law graph showing pressure is inversely proportional to volume when temperature is constant

• If the temperature increases, the graph is further from the origin and vice versa



IF THE VOLUME OF A GAS IS INCREASED, THE PARTICLES WILL BE FURTHER APART AND WILL COLLIDE LESS WITH EACH OTHER AND THE CONTAINER, DECREASING ITS PRESSURE

Copyright © Save My Exams. All Rights Reserved

Boyle's law: gas pressure is inversely proportional to the volume it occupies

Charles's Law

• If the pressure *P* of an ideal gas is constant, then **Charles's law** is given by:

 $V \propto T$

• This means the volume is **proportional** to the temperature of a gas

Page 12 of 33




• The relationship between the volume and thermodynamic temperature for a fixed mass of gas at constant pressure can also be written as:



 $\frac{V_1}{T_1} = \frac{V_2}{T_2}$

- Where:
 - V_1 = initial volume (m³)
 - V_2 = final volume (m³)
 - T_1 = initial temperature (K)
 - T_2 = final temperature (K)
- The variation of volume and temperature at a constant pressure is shown below:




Copyright © Save My Exams. All Rights Reserved

Charles's Law graph showing temperature (in K) is directly proportional to the volume when pressure is constant



Head to www.savemyexams.com for more awesome resources





IF THE TEMPERATURE OF A GAS IS INCREASED, THE PARTICLES GAIN KINETIC ENERGY AND MOVE FASTER. TO KEEP THE SAME PRESSURE, THEY MUST MOVE FURTHER APART WHICH INCREASES THE VOLUME OF THE GAS

Copyright © Save My Exams. All Rights Reserved

Charles's law: the temperature of a gas is proportional to the volume it occupies

Gay-Lussac's (Pressure) Law

■ If the volume V of an ideal gas is constant, the **Gay-Lussac's** or **Pressure law** is given by:

$$P \propto T$$


- This means the pressure is **proportional** to the temperature
- The relationship between the pressure and thermodynamic temperature for a fixed mass of gas at constant volume can also be written as:

$$\frac{P_1}{T_1} = \frac{P_2}{T_2}$$

- Where:
 - P_1 = initial pressure (Pa)
 - P_2 = final pressure (Pa)
 - T_1 = initial temperature (K)
 - T_2 = final temperature (K)
- The variation of pressure and temperature at a constant volume is shown below:




Head to www.savemyexams.com for more awesome resources



Copyright © Save My Exams. All Rights Reserved

TEMPERATURE (K)

Gay-Lussac's Law graph showing temperature (in K) is directly proportional to pressure when volume is constant



IF THE TEMPERATURE OF A GAS IS INCREASED, THE PARTICLES GAIN KINETIC ENERGY AND MOVE FASTER. THEREFORE THEY WILL COLLIDE MORE WITH EACH OTHER AND THE CONTAINER INCREASING ITS PRESSURE

Copyright © Save My Exams. All Rights Reserve

Pressure law: the temperature of a gas is proportional to the gas pressure

• Changes in the pressure, temperature or volume of an ideal gas are normally represented on a **pressure-volume** graph





• These are covered in more detail in the thermodynamics topic



Worked example

An ideal gas is in a container of volume 4.5×10^{-3} m³. The gas is at a temperature of 30 °C and a pressure of 6.2×10^5 Pa.

Calculate the pressure of the ideal gas in the same container when it is heated to 40 °C.

Answer:

Step 1: List the known quantities

- Volume, $V = 4.5 \times 10^{-3} \,\text{m}^3$
- Inital temperature, T₁ = 30 °C
- Pressure, $P = 6.2 \times 10^5 \, \text{Pa}$
- Final temperature, $T_2 = 40 \,^{\circ}\text{C}$

Step 2: State the ideal gas equation

$$pV = kT$$

• Where k = the constant of proportionality

Step 3: Rearrange for the constant of proportionality

$$k = \frac{pV}{T}$$

Step 4: Convert temperature Tinto Kelvin

$$\theta$$
 °C + 273.15 = TK

Step 5: Substitute in known value into the constant of the proportionality equation

$$k = \frac{6.2 \times 10^5 \times 4.5 \times 10^{-3}}{303.15} = 9.203...$$

Step 6: Rearrange the ideal gas relation equation for pressure

$$p = \frac{kT}{V}$$

Step 7: Substitute in new values

$$k = 9.203$$

$$T = 40 \,^{\circ}\text{C} = 40 + 273.15 = 313.15 \,\text{K}$$

Page 17 of 33



Head to www.savemyexams.com for more awesome resources

$$p = \frac{(9.203...) \times 313.15}{4.5 \times 10^{-3}} = 640.45... \times 10^{3} \text{ Pa} = 640 \text{ kPa}$$



Examiner Tip

Make sure to convert the temperature into **kelvin** before substituting into any gas law equation.

To know which gas laws to use, look at what values you've already been provided in the question. If it helps, label these as V_1 , T_1 etc. or look out for what quantity has been kept **constant**.

In the worked example, make sure not to get mixed up with this constant k with the Boltzmann constant $k_{\rm B}$. The constant depends on which ideal gas equation you use. It will be either nR or $k_{\rm B}$ T.

Ideal Gas Equation

Your notes

Ideal Gas Equation

• The two ideal gas equations, derived from the empirical gas laws, are:

$$PV = nRT$$

$$PV = Nk_BT$$

- The variables will be outlined below
- The empirical gas laws can be combined to give a single constant, known as the ideal gas constant, R

	Boyle's Law	Charles' Law	Pressure Law
relationship	PV = constant	$V \propto T$	$P \propto T$
constants	<i>T</i> , <i>n</i>	P, n	V, n

• An ideal gas is defined as:

A gas which obeys the ideal gas equation at all pressures, volumes and temperatures

• Combining the gas laws leads to the **ideal gas equation**:

$$PV = nRT$$

- Where:
 - P = pressure (Pa)
 - $V = \text{volume (m}^3)$
 - = n = number of moles (mol)
 - $R = 8.31 \,\mathrm{J}\,\mathrm{K}^{-1}\,\mathrm{mol}^{-1}$ (ideal gas constant)
 - T = temperature (K)

Constants

- The ideal gas constant R is the macroscopic equivalent of the Boltzmann constant $k_{_{I\!\!R}}$
 - The ideal gas constant is associated with macroscopic quantities such as volume and temperature
 - The Boltzmann constant is associated with the thermal energy of **microscopic** particles
- The **Boltzmann constant** is defined as the ratio of the ideal gas constant R and Avogadro's constant N_A :

$$k_B = \frac{R}{N_A}$$

- $\quad \text{Recall from the Amount of Substance revision note that } N_A \, = \, \frac{N}{n}$
- This gives another form of the ideal gas equation:

$$PV = Nk_BT$$

- Where:
 - \blacksquare N = number of molecules
 - $k_B = 1.38 \times 10^{-23} \text{ J K}^{-1}$ (Boltzmann constant)



Worked example

A gas has a temperature of -55° C and a pressure of 0.5 MPa. It occupies a volume of 0.02 m³.

Calculate the number of gas particles.

Answer:

Step 1: Write down the known quantities

- Temperature, *T* = -55°C = 218 K
- Pressure, $p = 0.5 \text{ MPa} = 0.5 \times 10^6 \text{ Pa}$
- Volume, V = 0.02 m³

Step 2: Write down the equation of state of ideal gases

$$PV = nRT$$

Step 3: Rearrange the above equation to calculate the number of moles n

$$n = \frac{PV}{RT}$$

Step 4: Substitute numbers into the equation

• From the data booklet, $R = 8.31 \,\mathrm{J \, K^{-1} \, mol^{-1}}$

$$n = \frac{(0.5 \times 10^6) \times 0.02}{8.31 \times 218} = 5.5 \text{ moles}$$

Step 5: Calculate the number of particles N

$$n = \frac{N}{N_A} \quad \Rightarrow \quad N = nN_A$$

$$N = 5.5 \times (6.02 \times 10^{23}) = 3.3 \times 10^{24}$$



Examiner Tip

The values for the gas constant, Avogadro constant and Boltzmann constant are all given in your data booklet.

Always make sure that temperature T is in **kelvin** for this topic. You must convert this from ${}^{\circ}$ C if not using the following conversion

$$\theta / {}^{\circ}C = T / K - 273$$

 $T / K = \theta / {}^{\circ}C + 273$





Head to www.savemyexams.com for more awesome resources

Kinetic Theory of Gases

Your notes

Kinetic Theory of Gases

- Ideal gases are described in terms of the kinetic theory
 - This is a modelled system used to approximate the behaviour of real gases
- A gas consists of atoms or molecules moving around randomly at high speeds
- The kinetic theory of gases models the thermodynamic behaviour of gases by linking:
 - The microscopic properties of particles e.g. mass and speed
 - The macroscopic properties of particles e.g. pressure and volume
- Kinetic theory and ideal gases are based on a set of the following assumptions:
 - 1. A gas consists of many identical molecules in a container. They all have the same mass
 - 2. The volume of the molecules is negligible **compared to** the volume of the container. This means they can be considered **point** particles
 - 3. The molecules are in continuous **random** motion at **high speeds**
 - 4. The molecules obey Newton's laws of motion
 - 5. The molecules collide elastically with each other and the walls of the container
 - 6. There are no intermolecular forces between the molecules **except during** collisions
 - 7. The time of a collision between molecules is negligible compared to the time between collisions
 - 8. External forces (e.g. gravity) are ignored
 - 9. The number of molecules of gas in a container is **very large**. Therefore the **average** behaviour (eg. speed) is usually considered
 - 10. Each particle exerts a force on the wall of the container with which it collides. This means the average of the forces produced by all gas particles results in a **uniform gas pressure**




Real & Ideal Gases

- Real gases will not always obey the ideal gas equation
- This is because the assumptions of the kinetic theory of gases may not always be valid
 - For example, the assumption that there are no intermolecular forces between gas molecules is not particularly realistic
- However, under certain conditions, they can be **considered** ideal gases
- An ideal gas is a good approximation of a real gas when:
 - The gas pressure is **low**
 - The gas density is low
 - Temperature is sufficiently higher than the boiling point of the substance
- At very high pressures, densities and low temperatures, real gases do not obey the kinetic theory model
- This is because at high pressures and densities:
 - Molecules are closer together
 - Therefore, there will be attractive forces between the molecules
 - The volume of the molecules cannot be considered negligible due to the high density
- At low temperatures:
 - A gas can change into a liquid, so the substance will no longer behave like a gas
- No gas is completely ideal, but air at normal room temperature and pressure, based on experiments, behaves very similar to an ideal gas







ying it a save my Exams. All highes heactived

Real vs. ideal gases





Head to www.savemyexams.com for more awesome resources

Derivation of the Kinetic Theory of Gases Equation


Your notes

Derivation of the Kinetic Theory of Gases Equation

- Gas pressure arises due to the collisions of the gas particles with the walls of the container that holds it
 - When a gas particle collides with a wall in the container, it exerts a **force** on the wall
 - The random motion of a large number of molecules exerting a force on the walls creates an overall **pressure**
 - This is because pressure = force per unit area

Derivation

- ullet Picture a single molecule in a cube-shaped box with sides of equal length I
- The molecule has a mass m and moves with speed c, parallel to one side of the box
- It collides at regular intervals with the ends of the box, exerting a force and contributing to the pressure of the gas
- By calculating the pressure this one molecule exerts on one end of the box, the total pressure produced by all the molecules can be deduced



A single molecule in a box collides with the walls and exerts a pressure

1. Find the change in momentum as a single molecule hits a wall perpendicularly

- One assumption of the kinetic theory is that molecules **rebound elastically**
 - This means there is no kinetic energy lost in the collision, so initial and final velocities are equal in magnitude
- When a gas particle of mass *m*, travels at an **average speed** *v*, and hits a wall of the container, it undergoes a **change in momentum** due to the force exerted by the wall on the particle


initial momentum = mV

final momentum = -mV

• Therefore, the change in momentum Δp is:

$$\Delta p$$
 = final momentum – initial momentum


$$\Delta p = -mv - (mv) = -2mv$$



$$F = \frac{\Delta p}{\Delta t} = -\frac{2mv}{\Delta t}$$

- According to Newton's Third Law, the particle exerts an equal and opposite force on the wall
- Therefore force exerted **on the wall** by the particle is:

$$F = \frac{2mv}{\Delta t}$$



A particle hitting a wall of the container in which the gas is held experiences a force from the wall and a change in momentum. The particle exerts an equal and opposite force on the wall

- 2. Calculate the number of collisions per second by the molecule on a wall
- The time between collisions from one wall to the other, and then back again, over a distance of 21 with speed v is:



time between collisions =
$$\frac{distance}{speed} = \frac{21}{v}$$



3. Find the change in momentum per second

• The force the molecule exerts on one wall, *F*, from step 1 now becomes the following when substituting the time between collisions for Δt:

$$F = \frac{2mv}{\Delta t} = \frac{2mv}{\frac{2l}{v}} = \frac{mv^2}{l}$$

4. Calculate the total pressure from N molecules

- The area of one wall is I^2
- The pressure, P can be written as:

$$P = \frac{F}{A} = \frac{\frac{mv^2}{l}}{l^2} = \frac{mv^2}{l^3}$$

- This is the pressure exerted from one molecule
- To account for the large number of N molecules, the pressure can now be written as:

$$P = \frac{Nmv^2}{l^3}$$

5. Consider the effect of the molecule moving in 3D space

- The pressure equation written above still assumes all the molecules are travelling in the same *x* direction and colliding with the same pair of opposite faces of the cube
 - To reflect this, it can be rewritten as:

$$P = \frac{Nmv_x^2}{l^3}$$

- Where v_x is the x component of the average velocity of all the particles
- In reality, all molecules will be moving in three dimensions equally
- Splitting the velocity into its components v_x , v_y and v_z to denote the amount in the x, y and z directions, v^2 (the square of the average speed) can be defined using pythagoras' theorem in 3D:

$$v^2 = v_x^2 + v_y^2 + v_z^2$$

• Since there is nothing special about any particular direction, it can be determined that the square of the average speed in each direction is equal:

$$v_x^2 = v_y^2 = v_z^2$$
$$v^2 = 3v_x^2$$



• Therefore, by combining the two equations, v can be defined as:

$$v_x^2 = \frac{1}{3}v^2$$

6. Re-write the pressure equation

- ullet The box is a cube with a side length I
- Therefore, volume of the cube $V=1^3$

$$P = \frac{Nmv^2}{3V}$$

• This can also be written using the density ρ of the gas:

$$\rho = \frac{mass}{volume} = \frac{Nm}{V}$$

- m is the mass of a single particle so Nm is the total mass of all the particles
- Rearranging the pressure equation for p and substituting the density ρ :

$$P = \frac{1}{3}\rho v^2$$

- Where:
 - P = pressure of the gas (Pa)
 - ρ = density of the gas (kg m⁻³)
 - $v = \text{mean square speed } (\text{m s}^{-1})$
- This is known as the **Kinetic theory of gases equation**

Worked example

An ideal gas has a density of 4.5 kg m^{-3} at a pressure of $9.3 \times 10^5 \text{ Pa}$ and a temperature of 504 K.

Determine the mean square speed of the gas molecules at 504 K.

Answer:

Step 1: List the known quantities

- Density of the gas, $\rho = 4.5 \text{ kg m}^{-3}$
- Gas pressure, $P = 9.3 \times 10^5 \, \text{Pa}$
- Temperature, $T = 504 \,\mathrm{K}$

Step 2: Rearrange the pressure equation for the mean square speed

$$P = \frac{1}{3}\rho v^2$$

$$v = \sqrt{\frac{3P}{\rho}}$$

Step 3: Substitute in the values

$$v = \sqrt{3 \times \frac{9.3 \times 10^5}{4.5}} = 787.4 = 790 \text{ m s}^{-1}$$



Examiner Tip

Although you will not be asked to recall the derivation, you need to understand the physics that underlines it and how to use the equation in the final result.

The 'mean square' speed just means an average speed, as we're assuming that all the molecules travel at the same speed.



Average Kinetic Energy of a Molecule

Your notes

Average Kinetic Energy of a Molecule

- We think of an ideal gas as molecules that collide elastically in random motion
- When the molecules collide, momentum and energy are conserved
 - We assume that, when the molecules are **not in contact**, no forces act between them
- This means they have no potential energy
 - Therefore, the internal energy *U* of an ideal, monatomic gas is equal to its **average kinetic energy** if the molecules are far enough apart
- The average kinetic energy, E_k for **one** molecule is equal to:

$$E_k = \frac{1}{2}mv^2 = \frac{3}{2}k_BT$$

- Where:
 - E_k = average kinetic energy of one molecule (J)
 - $v = \text{mean square speed of one molecule } (\text{m s}^{-1})$
 - m = mass of one molecule (kg)
 - k_B = Boltzmann constant
 - T = temperature of the gas (K)
- A consequence of this equation is that a greater gas temperature means a greater average kinetic
 energy of the particles
- Since the **total** internal energy is the **total** kinetic energy, the internal energy of a gas *U* is defined as:

$$U = \frac{3}{2} N k_B T$$

- Where:
 - U = internal energy (J)
 - N = number of molecules
- The relation to the amount of substance is:

$$U = \frac{3}{2}nRT$$


- Where:
 - = n = number of moles
 - \blacksquare R = ideal gas constant
- When heat is transferred to a fixed volume of gas, the internal energy increases and hence, so does the temperature, since the equations show that

$$U \propto T$$



Head to www.savemyexams.com for more awesome resources

- This is only relevant for **monatomic** gases. Examples include:
 - Helium
 - Neon
 - Argon
- A monatomic (one atom) molecule only has translational energy, whilst a diatomic (two-atom) molecule has both translational and rotational energy



A diatomic molecule has both rotational and translational kinetic energy



Worked example

600 J of thermal energy is transferred to 3 g of helium gas kept at a constant volume in a cylinder.

Helium has a mass number of 4.

Calculate the temperature of the gas.

Answer:

Step 1: List the known quantities

- Thermal (internal) energy, U = 600 J
- Molar mass of helium, $m_r = 4 \text{ g mol}^{-1}$ (from its mass number)
- Mass of helium gas, m = 3 g

Step 2: State the relevant equation

$$U = \frac{3}{2}nRT$$

Step 3: Calculate the number of moles of the gas

$$m_r = \frac{m}{n} \Rightarrow n = \frac{m}{m_r}$$

$$n = \frac{3}{4} = 0.75$$
 moles

Step 4: Rearrange the internal energy for the temperature, T

$$T = \frac{2U}{3nR}$$

Step 5: Substitute in the values

$$T = \frac{2 \times 600}{3 \times 0.75 \times 8.31} = 64 \,\mathrm{K}$$





Head to www.savemyexams.com for more awesome resources

Examiner Tip

Calculations involving moles, molar mass and number of molecules are common in this topic. You should be confident with converting between these quantities for all calculations in thermal physics.

Remember the combination 'nRT' and ' Nk_BT ' is also in the **ideal gas equation**. This shows how all the equations link together.

If you are using the equation $E_k=\frac{3}{2}k_BT$, remember this is only for **one** molecule, so you will need to multiply it by the total number of molecules for a whole gas.

