

# Practice Paper 2

Scan here to return to the course  
or visit [savemyexams.com](https://www.savemyexams.com)



---

Total Marks

/110

- 1 (a)** The following table shows the mean height,  $y$  cm, of primary school children who are age  $x$  years old.

<b>Age, <math>x</math> years</b>	6.25	7.35	8.5	9.25	10.75
<b>Mean Height, <math>y</math> cm</b>	115	121	129	136	140

The relationship between  $x$  and  $y$  can be modelled by the regression line of  $y$  on  $x$  with equation  $y = ax + b$ .

- i) Find the value of  $a$  and the value of  $b$ .
- ii) Write down the value of Pearson's product-moment correlation coefficient,  $r$ .

**(4 marks)**

- (b)** Use your regression equation from part (a) (i) to estimate the height of a child aged 9 years old.

**(2 marks)**

- (c)** Explain why it is not appropriate to use the regression equation to estimate the age of a child who is 133 cm tall.

**(1 mark)**

**2 (a)** An arithmetic sequence with a common difference  $-3.5$  has first term  $77$ .

Given that the  $r$ th term of the sequence is zero, find the value of  $r$ .

**(2 marks)**

**(b)** Find the maximum value of the sum of the first  $n$  terms of the sequence.

**(3 marks)**

**3**  $A$  and  $B$  are independent events, such that  $P(A) = 0.25$  and  $P(B) = 0.52$ .  $C$  is another event, such that  $B$  and  $C$  are mutually exclusive and  $P(A \cap C) = 0.09$ .

Given that  $P(A \cup B \cup C) = 0.95$ , find

i)  $P(A \cap B)$

ii)  $P(C)$

iii)  $P(A' \cap B')$

iv)  $P(A|C')$

(9 marks)

**4 (a)** Let  $f(x) = \frac{5-x^2}{3}$  and  $g(x) = 4 - \frac{3}{x}$ , where each function has the largest possible valid domain.

Write down the range of  $f$ .

**(1 mark)**

**(b)** Write down the domain and range of  $g$ .

**(2 marks)**

**(c)** Find

- i)  $(f \circ g)(x)$
- ii)  $(g \circ f)(x)$ .

**(3 marks)**

**5 (a)** The number of bacteria,  $n$ , in a dish, after  $t$  minutes is given by  $n = 5231e^{0.12t}$ .

Find the initial amount of bacteria.

**(2 marks)**

**(b)** Find the amount of bacteria after 12 minutes. Give your answer in the form  $a \times 10^k$ , where  $1 \leq a < 10$ ,  $k \in \mathbb{Z}$ .

**(3 marks)**

**(c)** Find the value of  $t$  when  $n = 2.7 \times 10^4$ .

**(2 marks)**

- 6 (a)** A UK energy company charges £0.22 per kilowatt hour (kWh) of electricity used. The amount of energy used per day by the company's customers,  $X$  kWh, follows the following probability density function

$$f(x) = \begin{cases} \frac{x(k-x)}{972}, & 0 \leq x \leq 18 \\ 0, & \text{otherwise} \end{cases}$$

Show that  $k = 18$ .

**(2 marks)**

- (b)** A customer's total daily charge consists of a fixed (standing) charge of £0.38 per day plus the charge for the electricity used.

- (i) Find the expected total daily charge.
- (ii) Find the standard deviation for the total daily charge.

**(6 marks)**

- 7** Consider the nine letters in the word MAGNITUDE.

Find the number of ways that the nine letters may be arranged if

- (i) there are no restrictions
- (ii) the four vowels (A, I, U, E) must all be together
- (iii) the arrangement starts with the letter M and ends with the letter E.

(5 marks)

8 Consider  $z = \text{cis } \theta$  where  $z \in \mathbb{C}$ ,  $z \neq 1$ .

Show that  $\text{Re}\left(\frac{1+z}{1-z}\right) = 0$ .

(5 marks)



**9 (a)** The binomial series expansion for  $(1 + t)^{-1}$  is given by

$$(1 + t)^{-1} = 1 - t + t^2 - \dots$$

Using the above result and the Maclaurin series for  $\cos(2x)$ , show that the Maclaurin series for  $\sec(2x)$  is

$$1 + 2x^2 + \frac{10}{3}x^4 + \dots$$

**(5 marks)**

**(b)** By using the result from part (a) and the Maclaurin series for  $\ln(1 + x)$ , find the value of the limit

$$\lim_{x \rightarrow 0} \left( \frac{x \ln(1 + 3x)}{\sec(2x) - 1} \right)$$

**(3 marks)**

- 10 (a)** The Strike A Light! matchstick company produces matchsticks with a length,  $X$  mm, that is normally distributed with mean 45 and variance  $\sigma^2$ .

The probability that  $X$  is greater than 45.37 is 0.1714.

Find  $P(44.63 < X < 45.37)$ .

**(2 marks)**

- (b)** i) Find  $\sigma$ , the standard deviation of  $X$ .  
ii) Hence, find the probability that a randomly selected matchstick has a length less than 44.5 mm.

**(5 marks)**

- (c)** Andrew has a box of Strike A Light! matches with fifteen matchsticks remaining in it. Those matchsticks may be assumed to be a random sample. Let  $Y$  represent the number of matchsticks in Andrew's box with lengths less than 44.5 mm.

Find  $E(Y)$ .

**(3 marks)**

- (d) Find the probability that exactly one of the matchsticks in Andrew's box has a length less than 44.5 mm.

**(2 marks)**

- (e) A Strike A Light! matchstick is selected at random and is found to have a length greater than 44.5 mm.

Find the probability that the length of the matchstick is between 44.63 mm and 45.37 mm.


**(3 marks)**


- 11 (a) Paola is modelling a small vase from her house for her maths project. To model the edge of the vase in cross-section, she decides to use a function  $f$  of the form

$$f(x) = \frac{qe^{\frac{x}{2}}}{2 + e^x}$$

where  $x \in \mathbb{R}$ ,  $x \geq 0$  and  $q \in \mathbb{R}^+$ .

The function and the vase are represented in the diagrams below.





The vertical height of the vase,  $OB$ , is measured along the  $x$ -axis. The radius of the vase's opening is  $OA$ , and its base radius is  $BC$ .

To model the vase, she will rotate by  $2\pi$  radians about the  $x$ -axis the region enclosed by the graph of  $y = f(x)$ , the  $x$ -axis, the  $y$ -axis, and the line  $x = \ln 43$ .

Show that the volume of the solid of revolution thus formed is  $\frac{14q^2\pi}{45}$  units<sup>3</sup>.

**(6 marks)**

**(b)** The volume of the actual vase is  $100 \text{ cm}^3$ .

Use this information to find the value of  $q$ .

**(2 marks)**

**(c)** Find the cross-sectional radius of the vase

(i) at its base,

(ii) at its widest point.

**(4 marks)**

**(d)** Paola wants to investigate how the cross-sectional radius of the vase changes.

Sketch a graph of the derivative of  $f$ , and use it to find the value of  $x$  at which the cross-sectional radius of the vase is decreasing most rapidly.

**(4 marks)**

**12 (a)** A function  $g$  is defined by  $g(x) = \arccos\left(\frac{x^2 - 1}{x^2 + 1}\right)$ ,  $x \in \mathbb{R}$ .

Show that  $g$  is an even function.

**(1 mark)**

**(b)** By considering the limit of  $g$  as  $x$  tends to infinity, show that the graph of  $y = g(x)$  has a horizontal asymptote and state its equation.

**(2 marks)**

**(c)** (i) Show that  $g'(x) = \frac{-2x}{(\sqrt{x^2})(x^2 + 1)}$  for  $x \in \mathbb{R}$ ,  $x \geq 0$ .

(ii) Considering the fact that  $\sqrt{x^2} = |x|$ , and also the expression for  $g'(x)$  above, show that  $g$  is increasing for  $x < 0$ .

**(9 marks)**

(d) A new function,  $h$ , is created by restricting the domain of  $g$ , such that

$$h(x) = \arccos\left(\frac{x^2 - 1}{x^2 + 1}\right), x \in \mathbb{R}, x \geq 0, \dots$$

Find an expression for  $h^{-1}(x)$ , carefully considering the range of  $h$  in determining your final answer.

(5 marks)

(e) State the domain of  $h^{-1}(x)$ .

(2 marks)