

# DP IB Maths: AA HL



## 2.9 Further Functions & Graphs

### Contents

- \* 2.9.1 Modulus Functions
- \* 2.9.2 Modulus Transformations
- \* 2.9.3 Modulus Equations & Inequalities
- \* 2.9.4 Reciprocal & Square Transformations



Your notes

## 2.9.1 Modulus Functions


### Modulus Functions & Graphs

#### What is the modulus function?

- The **modulus function** is defined by  $f(x) = |x|$ 
  - $|x| = \sqrt{x^2}$
  - Equivalently it can be defined  $|x| = \begin{cases} x & x \geq 0 \\ -x & x < 0 \end{cases}$
- Its **domain** is the set of **all real values**
- Its **range** is the set of **all real non-negative values**
- The modulus function gives the **distance** between 0 and  $x$ 
  - This is also called the **absolute value** of  $x$

#### What are the key features of the modulus graph: $y = |x|$ ?

- The graph has a **y-intercept** at  $(0, 0)$
- The graph has **one root** at  $(0, 0)$
- The graph has a **vertex** at  $(0, 0)$
- The graph is **symmetrical** about the **y-axis**
- At the **origin**
  - The function is **continuous**
  - The function is **not differentiable**



Copyright © Save My Exams. All Rights Reserved



Your notes

## What are the key features of the modulus graph: $y = a|x + p| + q$ ?

- Every **modulus graph** which is formed by **linear transformations** can be written in this form using key features of the modulus function
  - $|ax| = |a||x|$ 
    - For example:  $|2x + 1| = 2\left|x + \frac{1}{2}\right|$
  - $|p - x| = |x - p|$ 
    - For example:  $|4 - x| = |x - 4|$
- The graph has a **y-intercept** when  $x = 0$
- The graph can have 0, 1 or 2 **roots**
  - If  $a$  and  $q$  have the **same sign** then there will be **0 roots**
  - If  $q = 0$  then there will be **1 root** at  $(-p, 0)$
  - If  $a$  and  $q$  have **different signs** then there will be **2 roots** at  $\left(-p \pm \frac{q}{a}, 0\right)$
- The graph has a **vertex** at  $(-p, q)$
- The graph is **symmetrical** about the line  $x = -p$
- The value of  $a$  determines the **shape** and the **steepness** of the graph
  - If  $a$  is **positive** the graph looks like  $\nabla$
  - If  $a$  is **negative** the graph looks like  $\wedge$
  - The **larger** the value of  $|a|$  the **steeper** the lines
- At the **vertex**
  - The function is **continuous**
  - The function is **not differentiable**



Your notes

## 2.9.2 Modulus Transformations

### Modulus Transformations

#### How do I sketch the graph of the modulus of a function: $y = |f(x)|$ ?

- **STEP 1:** Keep the parts of the graph of  $y = f(x)$  that are **on or above the x-axis**
- **STEP 2:** Any parts of the **graph below the x-axis** get **reflected** in the x-axis

#### How do I sketch the graph of a function of a modulus: $y = f(|x|)$ ?

- **STEP 1:** Keep the graph of  $y = f(x)$  **only for  $x \geq 0$**
- **STEP 2:** **Reflect** this in the **y-axis**

#### What is the difference between $y = |f(x)|$ and $y = f(|x|)$ ?

- The graph of  $y = |f(x)|$  **never goes below the x-axis**
  - It does not have to have any lines of symmetry
- The graph of  $y = f(|x|)$  is **always symmetrical about the x-axis**
  - It can go below the x-axis

#### When multiple transformations are involved how do I determine the order?

- The transformations **outside the function** follow the **same order** as the **order of operations**
  - $y = |af(x) + b|$ 
    - Deal with the  $a$  then the  $b$  then the modulus
  - $y = a|f(x)| + b$ 
    - Deal with the modulus then the  $a$  then the  $b$
- The transformations **inside the function** are in the **reverse order** to the **order of operations**
  - $y = f(|ax + b|)$ 
    - Deal with the modulus then the  $b$  then the  $a$
  - $y = f(a|x| + b)$ 
    - Deal with the  $b$  then the  $a$  then the modulus

#### Examiner Tip


- When sketching one of these transformations in an exam question make sure that the graphs do not look smooth at the points where the original graph have been reflected
  - For  $y = |f(x)|$  the graph should look "sharp" at the points where it has been reflected on the x-axis
  - For  $y = f(|x|)$  the graph should look "sharp" at the point where it has been reflected on the y-axis



Your notes

 **Worked example**

The diagram below shows the graph of  $y = f(x)$ .



(a) Sketch the graph of  $y = |f(x)|$ .




Your notes

If the graph is on or above the  $x$ -axis then it stays the same

If the graph is below the  $x$ -axis then it is reflected in the  $x$ -axis

A stays the same  $(-1, 5)$

B becomes  $(3, 3)$




(b) Sketch the graph of  $y = f(|x|)$ .

keep the graph for  $x \geq 0$

Reflect this in the  $y$ -axis

A disappears

B stays the same  $(3, -3)$





Your notes

## 2.9.3 Modulus Equations & Inequalities

### Modulus Equations

#### How do I find the modulus of a function?

- The **modulus of a function**  $f(x)$  is
  - $|f(x)| = \begin{cases} f(x) & f(x) \geq 0 \\ -f(x) & f(x) < 0 \end{cases}$  or
  - $|f(x)| = \sqrt{[f(x)]^2}$

#### How do I solve modulus equations graphically?

- To solve  $|f(x)| = g(x)$  graphically
  - Draw  $y = |f(x)|$  and  $y = g(x)$  into your GDC
  - Find the  $x$ -coordinates of the **points of intersection**

#### How do I solve modulus equations analytically?

- To solve  $|f(x)| = g(x)$  analytically
  - Form **two equations**
    - $f(x) = g(x)$
    - $f(x) = -g(x)$
  - Solve both equations
  - **Check solutions** work in the original equation
    - For example:  $x - 2 = 2x - 3$  has solution  $x = 1$
    - But  $|(1) - 2| = 1$  and  $2(1) - 3 = -1$
    - So  $x = 1$  is not a solution to  $|x - 2| = 2x - 3$



Your notes

 **Worked example**

Solve for  $x$ :

a)  $\left| \frac{2x+3}{2-x} \right| = 5$

Analytically  
Split into two equations

$$\frac{2x+3}{2-x} = \pm 5$$

Solve individually

$$\frac{2x+3}{2-x} = 5$$

$$2x+3 = 10 - 5x$$

$$7x = 7$$

$$x = 1$$

$$\frac{2x+3}{2-x} = -5$$

$$2x+3 = 5x - 10$$

$$13 = 3x$$

$$x = \frac{13}{3}$$


Check:

$$\left| \frac{2(1)+3}{2-(1)} \right| = 5 \checkmark$$

$$\left| \frac{2(\frac{13}{3})+3}{2-(\frac{13}{3})} \right| = 5 \checkmark$$

$$x = 1 \text{ or } x = \frac{13}{3}$$

Graphically  
Sketch the two graphs



Find the points of intersection

$$(1, 5) \quad (4.33, 5)$$

Choose the  $x$ -coordinates

$$x = 1 \text{ or } x = 4.33 \text{ (3sf)}$$

b)  $|3x - 1| = 5x - 11$





Your notes

Analytically

Split into two equations

$$3x - 1 = \pm(5x - 11)$$

Solve individually

$$3x - 1 = 5x - 11$$

$$10 = 2x$$

$$x = 5$$

$$3x - 1 = 11 - 5x$$

$$8x = 12$$

$$x = 1.5$$

Check:

$$|3(5) - 1| = 14 \quad \checkmark$$

$$5(5) - 11 = 14 \quad \checkmark$$


$$|3(1.5) - 1| = 3.5$$

$$5(1.5) - 11 = -3.5 \quad \times$$

$$x = 5$$

Graphically

Sketch the two graphs



Find the points of intersection

$$(5, 14)$$

Choose the x-coordinates

$$x = 5$$

## Modulus Inequalities

### How do I solve modulus inequalities analytically?

- To solve **any** modulus inequality
  - First solve the corresponding modulus equation
    - Remembering to **check whether solutions are valid**
  - Then use a graphical method or a sign table to find the intervals that satisfy the inequality
- Another method is to solve **two pairs of inequalities**
  - For  $|f(x)| < g(x)$  solve:
    - $f(x) < g(x)$  when  $f(x) \geq 0$
    - $f(x) > -g(x)$  when  $f(x) \leq 0$
  - For  $|f(x)| > g(x)$  solve:
    - $f(x) > g(x)$  when  $f(x) \geq 0$
    - $f(x) < -g(x)$  when  $f(x) \leq 0$

#### Examiner Tip

- If a question on this appears on a calculator paper then use the same ideas as solving other inequalities
  - Sketch the graphs and find the intersections



Your notes



Your notes

### Worked example

 Solve the following inequalities for  $x$ .

a)  $|2x - 1| < 4$

 Solve for  $2x - 1 \geq 0$ 

For  $x \geq \frac{1}{2}$ :  $2x - 1 < 4 \Rightarrow x < \frac{5}{2} \therefore \frac{1}{2} \leq x < \frac{5}{2}$

 Solve for  $2x - 1 < 0$ 

For  $x < \frac{1}{2}$ :  $2x - 1 > -4 \Rightarrow x > -\frac{3}{2} \therefore -\frac{3}{2} < x < \frac{1}{2}$

Combine inequalities  $\boxed{-\frac{3}{2} < x < \frac{5}{2}}$

b)  $|x + 1| < |2x + 3|$

Solve the corresponding equation

$|x + 1| = |2x + 3| \Rightarrow x + 1 = \pm(2x + 3)$

Solve $x + 1 = 2x + 3$ $x = -2$	$x + 1 = -2x - 3$ $x = -\frac{4}{3}$
------------------------------------	---

Check $ (-2) + 1  = 1$ $ 2(-2) + 3  = 1$ ✓	$ (-\frac{4}{3}) + 1  = \frac{1}{3}$ ✓ $ 2(-\frac{4}{3}) + 3  = \frac{1}{3}$ ✓
---	---

Use a sign table

Check $x = -3$ $ (-3) + 1  <  2(-3) + 3 $ $2 < 3$ True ✓	Check $x = 1.5$ $ 1.5 + 1  <  2(1.5) + 3 $ $0.5 < 0$ False ✗	Check $x = 0$ $ 0 + 1  <  2(0) + 3 $ $1 < 3$ True ✓
--	--	---

Write solution  $\boxed{x < -2 \text{ or } x > -\frac{4}{3}}$



Your notes

## 2.9.4 Reciprocal & Square Transformations

### Reciprocal Transformations

#### What effects do reciprocal transformations have on the graphs?

- The **x-coordinates stay the same**
- The **y-coordinates change**
  - Their values become their **reciprocals**
- The coordinates  $(x, y)$  become  $\left(x, \frac{1}{y}\right)$  where  $y \neq 0$ 
  - If  $y = 0$  then a vertical asymptote goes through the original coordinate
  - Points that lie on the line  **$y = 1$**  or the line  **$y = -1$**  stay the same

#### How do I sketch the graph of the reciprocal of a function: $y = 1/f(x)$ ?

- Sketch the **reciprocal transformation** by considering the **different features** of the original graph
- Consider key points on the original graph
  - If  $(x_1, y_1)$  is a point on  $y = f(x)$  where  $y_1 \neq 0$ 
    - $\left(x_1, \frac{1}{y_1}\right)$  is a point on  $y = \frac{1}{f(x)}$ 
      - If  $|y_1| < 1$  then the point gets **further away from the x-axis**
      - If  $|y_1| > 1$  then the point gets **closer to the x-axis**
- If  $y = f(x)$  has a **y-intercept** at  $(0, c)$  where  $c \neq 0$ 
  - The reciprocal graph  $y = \frac{1}{f(x)}$  has a **y-intercept** at  $\left(0, \frac{1}{c}\right)$
- If  $y = f(x)$  has a **root** at  $(a, 0)$ 
  - The reciprocal graph  $y = \frac{1}{f(x)}$  has a **vertical asymptote** at  $x = a$
- If  $y = f(x)$  has a **vertical asymptote** at  $x = a$ 
  - The reciprocal graph  $y = \frac{1}{f(x)}$  has a **discontinuity** at  $(a, 0)$ 
    - The **discontinuity** will look like a **root**



Your notes


- If  $y = f(x)$  has a **local maximum** at  $(x_1, y_1)$  where  $y_1 \neq 0$ 
  - The reciprocal graph  $y = \frac{1}{f(x)}$  has a **local minimum** at  $\left(x_1, \frac{1}{y_1}\right)$
- If  $y = f(x)$  has a **local minimum** at  $(x_1, y_1)$  where  $y_1 \neq 0$ 
  - The reciprocal graph  $y = \frac{1}{f(x)}$  has a **local maximum** at  $\left(x_1, \frac{1}{y_1}\right)$
- Consider key regions on the original graph
  - If  $y = f(x)$  is **positive** then  $y = \frac{1}{f(x)}$  is **positive**
    - If  $y = f(x)$  is **negative** then  $y = \frac{1}{f(x)}$  is **negative**
  - If  $y = f(x)$  is **increasing** then  $y = \frac{1}{f(x)}$  is **decreasing**
    - If  $y = f(x)$  is **decreasing** then  $y = \frac{1}{f(x)}$  is **increasing**
  - If  $y = f(x)$  has a **horizontal asymptote** at  $y = k$ 
    - $y = \frac{1}{f(x)}$  has a **horizontal asymptote** at  $y = \frac{1}{k}$  if  $k \neq 0$
    - $y = \frac{1}{f(x)}$  **tends to  $\pm \infty$**  if  $k = 0$
  - If  $y = f(x)$  **tends to  $\pm \infty$**  as  $x$  tends to  $+\infty$  or  $-\infty$ 
    - $y = \frac{1}{f(x)}$  has a **horizontal asymptote** at  $y = 0$



Your notes

**Worked example**

The diagram below shows the graph of  $y = f(x)$  which has a local maximum at the point A.




Sketch the graph of  $y = \frac{1}{f(x)}$ .



Your notes

A becomes local minimum  $(-5, -\frac{1}{2})$   
 Vertical asymptote becomes root  $(2, 0)$   
 B becomes  $(0, -\frac{1}{4})$   
 C becomes vertical asymptote  $x=10$   
 Horizontal asymptote  $y=-1$  remains





Your notes

## Square Transformations

### What effects do square transformations have on the graphs?

- The effects are **similar to** the transformation  $y = |f(x)|$ 
  - The parts **below the x-axis are reflected**
  - The **vertical distance** between a point and the x-axis is **squared**
    - This has the effect of **smoothing the curve** at the x-axis
- $y = [f(x)]^2$  is **never below the x-axis**
- The **x-coordinates stay the same**
- The **y-coordinates change**
  - Their values are **squared**
- The coordinates  $(x, y)$  become  $(x, y^2)$ 
  - Points that lie on the **x-axis** or the line  **$y = 1$**  stay the same

### How do I sketch the graph of the square of a function: $y = [f(x)]^2$ ?

- Sketch the **square transformation** by considering the **different features** of the original graph
- Consider key points on the original graph
  - If  $(x_1, y_1)$  is a point on  $y = f(x)$ 
    - $(x_1, y_1^2)$  is a point on  $y = [f(x)]^2$ 
      - If  $|y_1| < 1$  then the point gets **closer to the x-axis**
      - If  $|y_1| > 1$  then the point gets **further away from the x-axis**
  - If  $y = f(x)$  has a **y-intercept** at  $(0, c)$ 
    - The square graph  $y = [f(x)]^2$  has a **y-intercept** at  $(0, c^2)$
  - If  $y = f(x)$  has a **root** at  $(a, 0)$ 
    - The square graph  $y = [f(x)]^2$  has a **root and turning point** at  $(a, 0)$
  - If  $y = f(x)$  has a **vertical asymptote** at  $X = a$ 
    - The square graph  $y = [f(x)]^2$  has a **vertical asymptote** at  $X = a$
  - If  $y = f(x)$  has a **local maximum** at  $(x_1, y_1)$ 
    - The square graph  $y = [f(x)]^2$  has a **local maximum** at  $(x_1, y_1^2)$  if  $y_1 > 0$
    - The square graph  $y = [f(x)]^2$  has a **local minimum** at  $(x_1, y_1^2)$  if  $y_1 < 0$
  - If  $y = f(x)$  has a **local minimum** at  $(x_1, y_1)$ 
    - The square graph  $y = [f(x)]^2$  has a **local minimum** at  $(x_1, y_1^2)$  if  $y_1 > 0$
    - The square graph  $y = [f(x)]^2$  has a **local maximum** at  $(x_1, y_1^2)$  if  $y_1 < 0$



### Examiner Tip

- In an exam question when sketching  $y = [f(x)]^2$  make it clear that the points where the new graph touches the x-axis are smooth
  - This will make it clear to the examiner that you understand the difference between the roots of the graphs  $y = |f(x)|$  and  $y = [f(x)]^2$




Your notes



Your notes

**Worked example**

The diagram below shows the graph of  $y = f(x)$  which has a local maximum at the point A.



Sketch the graph of  $y = [f(x)]^2$ .

- A becomes local minimum  $(-5, 4)$
- Vertical asymptote  $x = 2$  remains
- B becomes  $(0, 16)$
- C becomes local minimum
- Horizontal asymptote becomes  $y = 1$

