

DP IB Chemistry: HL



8.1 Theories & Reactions of Acids & Bases

Contents


- * 8.1.1 Brønsted-Lowry Acids & Bases
- * 8.1.2 Conjugate Acid-Base Pairs
- * 8.1.3 Characteristic Reactions of Acids
- * 8.1.4 Neutralization

8.1.1 Brønsted-Lowry Acids & Bases

Your notes


Brønsted-Lowry Acids & Bases

- The Brønsted-Lowry Theory defines acids and bases in terms of proton transfer between chemical compounds
- A Brønsted-Lowry acid is a species that gives away a proton (H+)
- A Brønsted-Lowry base is a species that accepts a proton (H+) using its lone pair of electrons



The diagram shows a Brønsted-Lowry acid which donates the proton to the Brønsted-Lowry base that accepts the proton using its lone pair of electrons

• The Brønsted-Lowry Theory is not limited to aqueous solutions only and can also be applied to reactions that occur in the gas phase



Page 2 of 15



Head to www.savemyexams.com for more awesome resources

Example of a Brønsted-Lowry acid and base reaction in the gas state



Worked example

Identify the correct role of the species in the following reaction:

$$H_2PO_4^-(aq) + H_2O(1) \rightarrow HPO_4^{2-}(aq) + H_3O^+(aq)$$

	Brønsted-Lowry Acid	Brønsted-Lowry Base
Α	H ₂ PO ₄ -	H ₂ O
В	H ₂ PO ₄ ²⁻	H ₂ PO ₄
С	H ₂ PO ₄	H ₃ O ⁺
D	H ₂ O	H ₂ PO ₄

Answer:

The correct option is **A**.

■ H₂PO₄⁻ is donating a proton to H₂O, so H₂PO₄⁻ must be an acid and H₂O must be a base

Examiner Tip

An atom of hydrogen contains 1 proton, 1 electron and 0 neutrons. When hydrogen loses an electron to become \mathbf{H}^+ only a **proton** remains, which is why a \mathbf{H}^+ ion is also called a proton.



8.1.2 Conjugate Acid-Base Pairs

Your notes

Conjugate Acid-Base Pairs

- A **Brønsted-Lowry acid** is a species that can donate a proton
- A **Brønsted-Lowry base** is a species that can accept a proton
- In a reaction at equilibrium, the products are formed at the same rate as the reactants are used

$$CH_3COOH(aq) + H_2O(I) = CH_3COO^-(aq) + H_3O^+(aq)$$

acid base conjugate base conjugate acid


- The reactant CH₃COOH is linked to the product CH₃COO⁻ by the transfer of a **proton** from the acid to the base
- Similarly, the H₂O molecule is linked to H₃O⁺ ion by the transfer of a proton
- These pairs are therefore called **conjugate acid-base pairs**
- A conjugate acid-base pair is two species that are different from each other by a H+ ion
 - Conjugate here means related
 - In other words, the acid and base are related to each other by one proton difference



Amphiprotic Species




- Species that can act both as proton donors and acceptors are called **amphiprotic**
 - Eg. water as a Brønsted-Lowry acid



The diagram shows water acting as a Brønsted-Lowry acid by donating a proton to ammonia which accepts the proton using its lone pair of electrons








Copyright © Save My Exams. All Rights Reserved

The Lewis diagram for the reaction of water with ammonia to show how water acts as a Brønsted-Lowry acid and ammonia as a Brønsted-Lowry base


■ Eg. water as a Brønsted-Lowry base







The diagram shows water acting as a Brønsted-Lowry base by accepting a proton from hydrochloric acid proton using its lone pair of electrons



The Lewis diagram for the reaction of water with hydrochloric acid to show how water acts as a Brønsted-Lowry base and ammonia as a Brønsted-Lowry acid



Head to www.savemyexams.com for more awesome resources

What is the difference between amphiprotic and amphoteric?

- A compound that is **amphoteric** means it has both basic and acidic character
 - When the compound reacts with an acid, it shows that it has basic character
 - When it reacts with a base, it shows that it's acidic
 - An example of this is aluminum oxide which reacts with both hydrochloric acid and sodium hydroxide:

$$Al_2O_3(s) + 6HCI(aq) \rightarrow 2AICI_3(aq) + 3H_2O(I)$$

 $Al_2O_3(s) + 2NaOH(aq) + 3H_2O(I) \rightarrow 2NaAI(OH)_4(aq)$

- When a compound is amphiprotic, it means it can act as a proton donor and as a proton acceptor
- Aluminium oxide is not amphiprotic, even though it is amphoteric

Amphiprotic versus Amphoteric Table

Amphiprotic	Amphoteric	
The term amphiprotic describes a substance that can both accept and donate a proton or H	The term amphoteric refers to the ability to act as both an acid and a base	
Amphiprotic substances can both accept or donate protons	Amphoteric substances can act as both an acid and a base	
All amphiprotic substances are amphoteric	Not all amphoteric substances are amphiprotic	

Copyright © Save My Exams. All Rights Reserved





Worked example

In the equilibrium reaction shown below, which species are a conjugate acid-base pair? $CH_3CH_2COOH(aq) + H_2O(I) = CH_3CH_2COO^{-}(aq) + H_3O^{+}(aq)$

- ${\bf A}$. CH₃CH₂CH₂COOH and H₂O
- \mathbf{B} . H_2O and H_3O^+
- C. H₂O and CH₃CH₂CH₂COO⁻
- D. CH₃CH₂CH₂COO⁻ and H₃O⁺



Answer

The correct option in **B**

■ A conjugate acid-base pair differ only by an H+ion

8.1.3 Characteristic Reactions of Acids

Your notes

Characteristic Reactions of Acids

Metals and acids

• The typical reaction of a metal and an acid can be summarized as

For example:

$$2HCl(aq) + Zn(s) \rightarrow ZnCl_2(aq) + H_2(g)$$

hydrochloric acid + zinc → zinc chloride + hydrogen

- Clearly, the extent of reaction depends on the **reactivity** of the metal and the **strength** of the acid
- Very reactive metals would react dangerously with acids and these reactions are not usually carried out
- Metals low in reactivity do not react at all, for instance copper does not react with dilute acids
- Stronger acids will react more vigorously with metals than weak acids. What signs of reaction would be expected to be different between the two?
 - Faster reaction, seen as
 - more effervescence
 - the metal dissolves faster
 - More exothermic

Metals and oxides

• The reaction of an acid with a metal oxide forms two products:

For example:

$$2HCI(aq) + CaO(s) \rightarrow CaCl_2(aq) + H_2O(l)$$

hydrochloric acid + calcium oxide → calcium chloride + water

Metals and hydroxides

• The reaction with a metal hydroxide and an acid follows the same pattern as an oxide:

• A suitable example might be:



$$H_2SO_4(aq) + Mg(OH)_2(s) \rightarrow MgSO_4(aq) + 2H_2O(l)$$

sulfuric acid + magnesium hydroxide → magnesium sulfate + water



Metals and carbonates

• The reaction between a metal carbonate and an acid produces three products:

For example:

$$2HNO_3(aq) + CuCO_3(s) \rightarrow Cu(NO_3)_2(aq) + H_2O(l) + CO_2(g)$$

nitric acid + copper carbonate → copper nitrate + water + carbon dioxide

Metals and hydrogencarbonates

• The reaction between a metal hydrogencarbonate and an acid is the same as the carbonate reaction with a slight difference in stoichiometry:

acid + metal hydrogencarbonate → salt + water + carbon dioxide

• An example of this would be:

$$HCI(aq) + NaHCO_3(s) \rightarrow NaCI(aq) + H_2O(l) + CO_2(g)$$

hydrochloric acid + sodium hydrogencarbonate → sodium chloride + water + carbon dioxide

Examiner Tip

Make sure you learn the formulae of the common acids and bases and that you can write examples of balanced equations of their characteristic reactions



Head to www.savemyexams.com for more awesome resources

Making Salts

- The acids and bases needed to make different salts can be deduced using the principles covered in the previous section
- The table below summarises these reactions

Making Salts Table

Type of salt	lon	Acid needed	Formula	Base needed
Sulfates	SO ₄ ²⁻	sulfuric	H ₂ SO ₄	
Nitrates	NO ₃	nitric	HNO ₃	metal oxide, hydroxide,
Chlorides	Cl⁻	hydrochloric	НСІ	carbonate or hydrogencarbonate
Ethanoates	CH³COO₋	ethanoic	CH₃COOH	
Ammonium	NH ₄	dny	_	aqueous ammonia

Copyright © Save My Exams. All Rights Reserved

• Note that although some metals can be used to make salts, they are not classified as bases as water is not a product of the reaction

Worked example

Which are the products of the reaction between zinc oxide and hydrochloric acid?

- A. zinc chloride and carbon dioxide
- B. zinc chloride, hydrogen gas and water
- C. zinc, hydrogen gas and water
- D. zinc chloride and water

Answer:

The correct option is ${\bf D}$.





• Metal oxides when reacting with acids produce a salt and water as the only products



8.1.4 Neutralization

Your notes

Neutralization

 A neutralisation reaction is one in which an acid (pH <7) and a base/alkali (pH >7) react together to form water (pH = 7) and a salt

• The proton of the acid reacts with the hydroxide of the base to form water

$$H^+(qq) + OH^-(qq) \xrightarrow{} H_2O(I)$$
Copyright © Save My Exams. All Rights Reserved

• The spectator ions which are not involved in the formation of water, form the salt

MAIN NEUTRALISATION REACTION:

THE TWO INDIVIDUAL REACTIONS TAKING PLACE ARE:

1.
$$H^+ + OH^- \longrightarrow H_2O$$

2. $Na^+ + Cl^- \longrightarrow NaCl$

The diagram shows a neutralisation reaction of HCl and NaOH and the two individual reactions that take place to form the water and salt

• The name of the salt produced can be predicted from the acid that has reacted

Acid Reacted & Salt Table



Acid Reacted	Salt produced		
Hydrochloric Acid	A Chloride		
Nitric Acid	A Nitrate		
Sulfuric Acid	A Sulfate		



Copyright © Save My Exams. All Rights Reserved



Examiner Tip

The enthalpy of neutralisation is the enthalpy change that occurs when an acid reacts with a base to form one mole of water. Since the reaction between strong acids and strong bases is the same regardless of the acid or base, it should be no surprise the enthalpy change is the same and is approximately -57 kJ mol⁻¹